

Research Journal of Mathematical and Statistical Sciences . Vol. 1(3), 10-15, April (2013)

A Study of Hall Currents on Magneto-Hydrodynamic Unsteady Flow of Visco-Elastic [Oldroyd (1958) Model] Fluid through Porous Media in a Rectangular Channel

Sharma A. K. ¹, Kumar Ravindra ² and Mishra Gaurav² ¹Department of Mathematics, K.K.P.G.College Etawah, U.P., INDIA ²Department of Mathematics, Agra College Agra, U.P., INDIA

Available online at: www.isca.in

Received 13th December 2012, revised 22nd February 2013, accepted 28th March 2013

Abstract

The object of the present paper is to study of Hall currents on magneto-hydrodynamic unsteady flow of visco-elastic first order Oldroyd model fluid with transient pressure gradient through porous media in a long rectangular channel. The expression for velocity of the fluid is obtained in elegant form. Some deductions have been discussed in detail.

Keywords: MHD flow, visco-elastic fluid, transient pressure gradient, porous media and hall current. msc 2000: 76a05, 76s05, 76s05

Introduction

The fluids which exhibit the elasticity property of solids and viscous property of liquids are called visco-elastic fluids or non-Newtonian fluids. In hydromagnetic flow we study of the flow of electrically conducting fluid in presence of Maxwell electromagnetic field. The flow of the conducting fluid is effectively changed by the presence of the magnetic field and the magnetic field is also perturbed due to the motion of the conducting fluid. This phenomena is therefore interlocking in character and the discipline of this branch of science is called Magnetohydrodynamics (MHD). It is equally rich and admits wider applications in engineering, technology, cusmology, Astrophysics and other applied sciences. It has tremendously developed in last forty years and some of the monographs in this field are due to Oldroyd¹, Ferraro and Plumton², Pai³, Shercliff⁴, Sutton and Shermann⁵, Jefferey⁶ and Cowling⁷.

Flow behaviour of visco-elastic fluids through channels of different cross-section have been studied by a number of authors Ghosh and Sengupta⁸, Kundu and Sengupta⁹, Abdul Hadi and Sharma¹⁰ and Mishra and Panda¹¹. Many research workers have paid their attention towards the application of visco-elastic fluid flow through various types of channel under the influence of magnetic field such as Sengupta and Basak¹², Kundu and Sengupta¹³, Sengupta and Paul¹⁴, Rehman and Alam Sarkar¹⁵, Krishna and Rao¹⁶, Ghosh and Ghosh¹⁷, Radhakrishnamacharya and Rao¹⁸ and Kumar, Singh and Sharma¹⁹.

The study of physics of flow through porous medium has become the basis of many scientific and engineering applications. This type of flow is of great importance in the petroleum engineering concerned with the movement of oil, gas and water through reservoir of oil or gas field to the hydrologist in the study of the migration of underground water and to the chemical engineer in the filtration process. Many research workers have paid their attention towards the application of visco-elastic fluid flow of different category through porous medium in chemicals of various cross-section such as Sudhakar and Venkataramana²⁰, Kumar and Singh²¹,Gupta and Gupta²², Hassanien²³, Agarwal and Agarwal²⁴, Sharma and Pareek²⁵, Ahmed²⁶, Thakur and Kumar²⁷, Singh, Kumar and Sharma²⁸, Singh, Mishra and Sharma²⁹, Das and Haque³⁰, Kumar, Sharma and Singh³¹ and Kumar, Mishra and Singh³² etc.

The object of the present paper is to study of Hall currents on magneto-hydrodynamic unsteady flow of visco-elastic first order Oldroyd¹ model fluid with transient pressure gradient through porous media in a long rectangular channel. The expression for velocity of the fluid is obtained in elegant form. Some deductions have been discussed in detail. In this paper we have considered the problem of Singh, Verma and Varshney³³ with porous media.

Basic Theory

For slow motion, the Rheological equations for Oldroyd visco-elastic liquid are:

Research Journal of Mathematical and Statistical Sciences _ Vol. 1(3), 10-15, April (2013)

ISSN 2320–6047 Res. J. Mathematical & Statistical Sci.

$$\tau_{ij} = -p \,\delta_{ij} + \tau'_{ij}$$

$$\left(1 + \lambda_1 \frac{\partial}{\partial t}\right) \tau'_{ij} = 2\mu \left(1 + \mu_1 \frac{\partial}{\partial t}\right) e_{ij}$$

$$e_{ij} = \frac{1}{2} \left(v_{i,j} + v_{j,i}\right)$$

$$(3)$$

$$v_{i,i} = 0$$
 (4)
where τ_{ij} = The stress tensor, τ'_{ij} = The deviatoric stress tensor, e_{ij} = The rate of strain tensor, p = The pressure, λ_1 = The

stress relaxation time parameter, μ_1 = The strain rate retardation time, δ_{ij} = The metric tensor, μ = The coefficient of viscosity, v_i = The velocity components

Formulation of the Problem

Using a rectangular Cartesian coordinate system (x, y, z) such that the z- axis is along the axis of the channel and the walls of the channel are taken to be planes $x = \pm a$ and $y = \pm b$. Let us consider the flow of visco-elastic Oldroyd liquid along the axis of rectangular channel i.e. along z-axis only, therefore 0, 0, w(x, y, z) are the velocity components along x, y, z directions respectively. A transient pressure gradient $-Pe^{-\omega t}$ varying with time *t* is applied to the liquid.

Following the stress-strain relations (1)-(4), the equation for unsteady motion through porous medium in a rectangular channel under the influence of an uniform magnetic field B_0 applied perpendicular to flow of the conducting visco-elastic Oldroyd liquid when induced magnetic field be neglected is given by

$$\left(1+\lambda_{1}\frac{\partial}{\partial t}\right)\frac{\partial w}{\partial t} = -\frac{1}{\rho}\left(1+\lambda_{1}\frac{\partial}{\partial t}\right)\frac{\partial p}{\partial z} + \nu\left(1+\mu_{1}\frac{\partial}{\partial t}\right)\left(\frac{\partial^{2}w}{\partial x^{2}} + \frac{\partial^{2}w}{\partial y^{2}}\right) - \left(1+\lambda_{1}\frac{\partial}{\partial t}\right)\frac{\sigma B_{0}^{2}}{\rho(1+m_{1}^{2})}w - \frac{\nu}{k}\left(1+\lambda_{1}\frac{\partial}{\partial t}\right)w \quad (5)$$

Where B_0 is an uniform magnetic field, σ the electrical conductivity, v the kinematic coefficient of viscosity, ρ the density, μ the coefficient of viscosity, m_1 Hall parameter and k is the permeability of porous medium.

Introducing the following non-dimensional quantities:

$$w^{*} = \frac{a}{v}w, \quad p^{*} = p\frac{a^{2}}{\rho v^{2}}, \quad t^{*} = \frac{v}{a^{2}}t, \quad \left(x^{*}, y^{*}, z^{*}\right) = \frac{1}{a}\left(x, y, z\right), \quad \omega^{*} = \omega\frac{a^{2}}{v}, \quad \lambda_{1}^{*} = \lambda_{1}\frac{v}{a^{2}}, \quad \mu_{1}^{*} = \frac{v}{a^{2}}\mu_{1}, \quad k^{*} = \frac{1}{a^{2}}k$$

in equation (1), we get (after dropping the stars)

$$\left(1 + \lambda_1 \frac{\partial}{\partial t}\right) \frac{\partial w}{\partial t} = -\left(1 + \lambda_1 \frac{\partial}{\partial t}\right) \frac{\partial p}{\partial z} + \left(1 + \mu_1 \frac{\partial}{\partial t}\right) \left(\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}\right) - \left(\frac{M^2}{(1 + m_1^2)} + \frac{1}{k}\right) \left(1 + \lambda_1 \frac{\partial}{\partial t}\right) w$$

$$(6)$$

where $M = aB_0 \sqrt{\frac{\sigma}{\mu}}$ (Hartmann number)

We have considered those types of situations of the flow which is transient in nature with respect to time and periodic in nature with respect to y. From the nature of the boundary conditions it is reasonable to choose the solution of (6) as

$$w = W(x)\cos my e^{-\omega t}$$
⁽⁷⁾

The boundary conditions are:

(i) W=0 when
$$x = \pm 1$$
, $-\frac{b}{a} \le y \le \frac{b}{a}$ (8)

International Science Congress Association

11

Research Journal of Mathematical and Statistical Sciences Vol. 1(3), 10-15, April (2013)

(ii) W=0 when
$$y = \pm \frac{b}{a}$$
, $-1 \le x \le 1$

Boundary condition (9) will be satisfied if $\cos m \frac{b}{a} = 0$

or
$$m \frac{b}{a} = (2n+1)\frac{\pi}{2}$$

or $m = (2n+1)\frac{\pi a}{2b}, \quad n = 0,1,2,3,....$ (10)

We construct the solution as the sum of all possible solutions for each value of n:

$$w = \sum_{n=0}^{\infty} W(x) \cos m \, y.e^{-\omega t} \tag{11}$$

By putting $\frac{\partial p}{\partial z} = -Pe^{-\omega t}$, $(\omega > 0)$ in (6) and using (11), we get

$$\sum_{n=0}^{\infty} \left\{ \frac{d^2 W(x)}{dx^2} - m^2 W(x) \right\} \cos my + \sum_{n=0}^{\infty} \frac{\omega(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} W(x) \cos my - \sum_{n=0}^{\infty} \frac{\left(\frac{M^2}{(1 + m_1^2)} + \frac{1}{k}\right)(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} \times W(x) \cos my + \frac{4P}{\pi} \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)} \cos my = 0$$

$$\sum_{n=0}^{\infty} \left[\left\{ \frac{d^2 W(x)}{dx^2} - \left(m^2 - \frac{\omega(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} + \left(\frac{M^2}{(1 + m_1^2)} + \frac{1}{k}\right) \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)}\right) W(x) + \frac{4P}{\pi} \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} \frac{(-1)^n}{(2n+1)} \right\} \cos my \right] = 0$$
(12)

Now from (12) equating the coefficient of cos my equal to zero, the value of W(x) can be determined from

$$\frac{d^2 W(x)}{dx^2} - \left\{ m^2 - \frac{\omega(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} + \left(\frac{M^2}{(1 + m_1^2)} + \frac{1}{k} \right) \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} \right\} W(x) + \frac{(-1)^n}{(2n+1)} \cdot \frac{4P}{\pi} \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} = 0$$
or
$$\frac{d^2 W(x)}{dx^2} - \frac{K^2}{a^2} W(x) + A_n = 0$$
(13)

where

$$K^{2} = \left\{ m^{2} - \frac{\omega(1 - \lambda_{1}\omega)}{(1 - \mu_{1}\omega)} + \left(\frac{M^{2}}{(1 + m_{1}^{2})} + \frac{1}{k} \right) \frac{(1 - \lambda_{1}\omega)}{(1 - \mu_{1}\omega)} \right\} a^{2} \text{ and } A_{n} = \frac{(-1)^{n}}{(2n+1)} \cdot \frac{4P}{\pi} \frac{(1 - \lambda_{1}\omega)}{(1 - \mu_{1}\omega)}$$
(14)

Solving (13) subject to boundary condition (8), we get

$$W(x) = \frac{A_n}{K^2} \left\{ 1 - \frac{\cos h \frac{K}{x}}{\cos h \frac{K}{a}} \right\}$$

or
$$W(x) = \frac{(-1)^n}{(2n+1)} \cdot \frac{4P}{\pi} \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega) K^2} \left\{ 1 - \frac{\cos h \frac{K}{a} x}{\cos h \frac{K}{a}} \right\}$$
(15)

Putting the value of W(x) in (11), we get the velocity of visco-elastic Oldroyd liquid under the influence of uniform transverse magnetic field.

ISSN 2320-6047 Res. J. Mathematical & Statistical Sci.

(9)

Research Journal of Mathematical and Statistical Sciences ____ Vol. 1(3), 10-15, April (2013)

ISSN 2320-6047 Res. J. Mathematical & Statistical Sci.

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^n}{(2n+1)} \cdot \frac{4P}{\pi} \frac{(1-\lambda_1 \omega)}{(1-\mu_1 \omega) K^2} \left\{ 1 - \frac{\cos h \frac{K}{a} x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(16)

Deductions

(i) Taking limit $\mu_1 \rightarrow 0$, the visco-elastic liquid becomes Maxwell liquid, the velocity is given by the expression:

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^n}{(2n+1)} \cdot \frac{4P}{\pi} \frac{(1-\lambda_1 \omega)}{K^2} \left\{ 1 - \frac{\cos h \frac{K}{a} x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(17)
where $K^2 = \left\{ (m^2 + \frac{M^2}{(1+m_{\star}^2)} + \frac{1}{k} - \omega)(1-\lambda_1 \omega) \right\} a^2$

 $\begin{pmatrix} (1+m_1^2) & k \end{pmatrix}$ (ii) Taking limit $\lambda_1 \to 0$, the visco-elastic liquid becomes Kuvshinisky liquid, the velocity is given by the expression:

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^{n} 4P}{(2n+1)\pi (1-\mu_{1}\omega)K^{2}} \left\{ 1 - \frac{\cos h \frac{K}{a}x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(18)
where
$$K^{2} = \left\{ m^{2} + \frac{\left(\frac{M^{2}}{(1+m_{1}^{2})} + \frac{1}{k} - \omega\right)}{(1-\mu_{1}\omega)} \right\} a^{2}$$

(iii) Taking limits $\mu_1 \rightarrow 0$ and $\lambda_1 \rightarrow 0$, the visco-elastic liquid becomes purely viscous liquid, the velocity is given by the expression: ~

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^{n} 4P}{(2n+1)\pi K^{2}} \left\{ 1 - \frac{\cos h \frac{K}{a} x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(19)
where $K^{2} = \left\{ m^{2} + \frac{M^{2}}{(1+m_{1}^{2})} + \frac{1}{k} - \omega \right\} a^{2}$

Particular cases

Case I: Taking limit $M \rightarrow 0$ i.e. if uniform magnetic field is withdrawn, the velocity of visco-elastic Oldroyd liquid through porous medium is given by the expression: Г ٦

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^n 4P(1-\lambda_1 \omega)}{(2n+1)\pi (1-\mu_1 \omega) K^2} \left\{ 1 - \frac{\cos h \frac{K}{-x}}{\cos h \frac{K}{-a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(20)

(

Research Journal of Mathematical and Statistical Sciences ______ Vol. 1(3), 10-15, April (2013) _____ ISSN 2320–6047 Res. J. Mathematical & Statistical Sci.

where
$$K^2 = \left\{ m^2 - \frac{\omega(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} - \frac{1}{k} \frac{(1 - \lambda_1 \omega)}{(1 - \mu_1 \omega)} \right\} a^2$$

Case II: Taking limit $k \to \infty$ i.e. porous medium is withdrawn, velocity of visco-elastic Oldroyd liquid under the influence of magnetic field is given by

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^{n} 4P(1 - \lambda_{1}\omega)}{(2n+1)\pi(1 - \mu_{1}\omega)K^{2}} \left\{ 1 - \frac{\cos h \frac{K}{a}x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(21)
where $K^{2} = \left\{ m^{2} - \frac{\omega(1 - \lambda_{1}\omega)}{(1 - \mu_{1}\omega)} + \frac{M^{2}}{(1 + m_{1}^{2})} - \frac{(1 - \lambda_{1}\omega)}{(1 - \mu_{1}\omega)} \right\} a^{2}$

Case III: Taking limit $M \to 0$ and $k \to \infty$ i.e. uniform magnetic field and porous media both are withdrawn, the velocity of visco-elastic Oldroyd liquid is given by

$$w = \sum_{n=0}^{\infty} \left[\frac{(-1)^{n} 4P(1-\lambda_{1}\omega)}{(2n+1)\pi(1-\mu_{1}\omega)K^{2}} \left\{ 1 - \frac{\cos h \frac{K}{a}x}{\cos h \frac{K}{a}} \right\} \cos \frac{(2n+1)\pi a}{2b} y \right] e^{-\omega t}$$
(22)
where $K^{2} = \left\{ m^{2} - \frac{\omega(1-\lambda_{1}\omega)}{(1-\mu_{1}\omega)} \right\} a^{2}$

All the expressions of Kundu and Sengupta⁹ are obtained.

Case IV: Putting $m_1 = 0$, we shall get the same result as Singh, Verma and Varshney³³.

Conclusion

The nature of the Magneto-hydrodynamics and porous medium is to reduce the velocity of the fluid therefore the presence of the porous medium in the rectangular channel under the influence of uniform magnetic field will definitely reduced the velocity of visco-elastic fluid and evidently velocity of the fluid in deductions case I, case II and case III will be slower than the velocity of the fluid obtained by Kundu and Sengupta⁹.

Acknowledgement

The authors are thankful to Dr. K. K. Singh, Ex. Reader, Department of Mathematics, Agra College, Agra (U.P.) for his valuable suggestions and moral support.

References

- 1. Oldroyd J.G., Proc. Roy. Soc. (London) 254 278 (1958)
- 2. Ferraro V.C.A. and Plumpton C., An Introduction to Magnetohydrodynamics, Oxford University Press (1961)
- 3. Pai S.I., Magnetohydrodynamics and Plasma dynamics, Prantice Hall London (1962)
- 4. Shercliff J.A., A Text Book of Magnetohydrodynamics, Pergamon Press (1965)
- 5. Sutton G.W. and Shermann A., Engg. Magnetohydrodynamics, McGraw Hill Book Company (1965)
- 6. Jefferey A., Magnetohydrodynamics, Oliver and Boyd., (1966)
- 7. Cowling T.G., Magnetohydrodynamics, Bristol, England Adam Higher Limited (1976)
- 8. Ghosh B.C. and Sengupta P.R., Proc. Nat. Acad. Sci. India 66A(1), 81 (1996)
- 9. Kundu S.K. and Sengupta P.R., Proc. Nat. Acad. Sci. India, 71(A)III 253 (2001)

Research Journal of Mathematical and Statistical Sciences _ Vol. 1(3), 10-15, April (2013)

- 10. Abdul Hadi A.M. and Sharma K.C., Ind. Jour. Pure Appl. Math. 33(1), 131 (2002)
- 11. Mishra B.K. and Panda T.C., Acta Ciencia Indica, XXXI(2M), 341 (2005)
- 12. Sengupta P.R. and Basak P., Ind. Jour. Theo. Phy., 50(3), 203 (2002)
- 13. Kundu S.K. and Sengupta P.R., Bull. of Allahabad Mathematical Soc., 18, 73 (2003)
- 14. Sengupta P.R. and Paul S., Ind. Jour. Theo. Phy. 52(4), 327 (2004)
- 15. Rehman M.M. and Alam Sarkar M.S., Bull. Cal. Math. Soc., 96(6), 463 (2004)
- 16. Krishna D.V. and Rao P.N., Acta Ciencia Indica, XXXI(3M), 917 (2005)
- 17. Ghosh, S. and Ghosh, A.K. Ind. Jour. Pure Appl. Math., 36(10), 529 (2006)
- 18. Radhakrishnamacharya G. and Rao S., Proc. Nat. Acad. Sci. India, 76 (A)III, 241 (2007)
- 19. Kumar R., Singh K.K. and Sharma A.K., Jour. Purvanchal Acad. Sci., 15, 80 (2009)
- 20. Sudhakar B. and Venkataramana S., Reg. Engg. Heat Mass Transfer, 10(3), 221 (1988)
- 21. Kumar G. and Singh K.K., Pure Appl. Mathematika Sci., XXXII(1-2), 65 (1990)
- 22. Gupta R.K. and Gupta K., Ind. Jour. Pure Appl. Maths., 27(4), 423 (1996)
- 23. Hassanien I.A. ZAMM, 82(6), 409 (2002)
- 24. Agarwal M. and Agarwal N., Proc. Nat. Acad. Sci. India, 76A(1), 55 (2006)
- 25. Sharma P.R. and Pareek D., Bull. Pure Appl. Sci., 20(1), 103 (2006)
- 26. Ahmed, S. Bull. Cal. Math. Soc., 99(5), 511 (2007)
- 27. Thakur C. and Kumar M., Pure Appl. Math. Sci., LXVII(1-2), 113 (2008)
- 28. Singh K.K, Kumar R. and Sharma A.K., Acta Ciencia Indica, XXXIV(1M), 351 (2008)
- 29. Singh K.K, Mishra G. and Sharma A.K., Acta Ciencia Indica, XXXIV(1), 421 (2008)
- 30. Das A. and Haque A., Ultra Scientists, 20(2M), 291 (2008)
- 31. Kumar R., Sharma A.K. and Singh K.K., Ultra Scientists, 20(2M), 545 (2008)
- 32. Kumar R., Mishra G. and Singh K.K., Int. Jour. Math. Arch., 2(4), 584 (2011)
- 33. Singh U.R., Verma D.K. and Varshney N.K., Int. Jour. Math. Arch., 3(8), 3216 (2012)