



Approximate Analysis of A Single Server Queue with Unit Step Function

Dhanesh Garg

Department of Mathematics, Maharishi Markandeshwar University, Mullana-133203, INDIA

Available online at: www.isca.in, www.isca.me

Received 24th July 2014, revised 11th September 2014, accepted 5th October 2014

Abstract

This study analyzed a single server queueing model with a time dependent arrival rate $\lambda(1+tT^{-1})$ and service rate μ is constant. In this model, the incoming arrivals are Poisson stream, service times are negative exponentially distributed and the first come first served queueing discipline. We obtain an explicit expression for the state probability distribution with time dependent arrival rate using unit step function. For a special form of the traffic intensity, If the service rate is constant ($=\mu$), this corresponds to $\rho(t) = \lambda(1+tT^{-1})\mu^{-1}$

Keywords: Transient analysis, Unit step function, Queueing system. AMS Subject Classification: 60K25, 90B25

Introduction

The vast majority of the queueing literature considers only models where the arrivals and service processes are time-homogeneous. Yet it is clear that in many situations it is quite natural to have, say, the arrival rate varying with time. Last some years a number of interesting literature that discuss the transient behaviour of the M/M/1 queue, an aspect which is very important for practical applications have focused by cf. Abate and Whitt¹⁻⁴. Transient solutions of queues have been obtained different methods: generating function method by Bailey⁵, spectral method by Lederman and Reuter⁶, difference equation method by Conolly⁷, alternative approach of generating functions by Parthasarathy⁸. Specific case studies and applications of time-dependent (i.e. non-stationary) queues are given in Clarke A.B. and Koopman B.O.⁹⁻¹⁰. The relatively small body of literature on time-dependent queues is probably due to the fact that standard methods for treating time-independent models, such as generating functions and Laplace transforms, do not easily generalize to non-stationary models. In most of the literature it is assumed that the parameters of the system are constant, i.e. do not vary over time. But in our model one parameter is time dependent and other is time independent. Our primary objective in this paper is to approximate the analysis of a single server queue, arrival rate varying with time and service rate is constant using step function. In this model, the incoming arrivals are Poisson stream, service times are negative exponentially distributed and the first come first served queueing discipline. For a special form of the traffic intensity, we obtain an explicit analytic expression for the probability distribution function with time dependent arrival rate using unit step function. The rest of the paper is organized as

follows: In the next section, the mathematical description of the considered queueing model. In section 3, the transient solution to the model derived.

Model description

We consider a single server queue at which customers arrive to a Poisson stream of rate. The service time is exponentially distributed time-independent rate. Let the system take initially non-empty with FCFS discipline i.e., "j" is in the system at t=0. In view of the arbitrary rate functions, there is no loss of generality in assuming that we start the process at t=0, by introduce a new function i.e. unit step function define as

$$1-u(t) = \begin{cases} 1 & \text{if } t < T \\ 0 & \text{if } t > T \end{cases} \quad (1)$$

This function if useful in expressing boundary conditions which is depends upon time. The basis of the approximate differential equation derived from the time dependent arrival rate and time independent service rate. Let $P_n(t)$ = the probability that there are n customers in the system at time t given that there were j customers in the system at time 0. Therefore,

Probability that [one arrival in the time interval $(t, t + \Delta t)$] = $\lambda(1+tT^{-1})(1-u_T(t))\Delta t + 0(\Delta t)$

Probability that [one departure in the time interval $(t, t + \Delta t)$] = $\mu\Delta t + 0(\Delta t)$

Transient state equations for the finite source M/M/1 queueing system are given by

$$\begin{aligned} \frac{d}{dt} p_n(t) &= -[\lambda(1+tT^{-1})(1-u_T(t)+\mu)] p_n(t) + \\ &[\lambda(1+tT^{-1})(1-u_T(t))] p_{n-1}(t) + \mu p_{n+1}(t) : n > 0 \quad (2-3) \\ \frac{d}{dt} p_0(t) &= -[\lambda(1+tT^{-1})(1-u_T(t)+\mu)] p_0(t) + \mu p_1(t) : n = 0 \end{aligned}$$

The model solution

The solution of our problem of this queueing system of differential-difference equations under the initial conditions

$P_n(t=0) = \psi_{jn}$, where ψ is the Kronecker delta

$$\psi_{jn} = \begin{cases} 1 & \text{if } n = j \\ 0 & \text{if } n \neq j \end{cases} \quad (4)$$

We define,

$$\frac{d}{dt} p_n(t) = \mu \frac{d}{d\pi} p_n(\pi)$$

$$Q_n(\pi, T) = \text{Exp.}\{\pi[1 + R(\pi, T)]\} p_n(\pi) \quad (6)$$

$$\rho(\pi, T) = \left[\frac{\lambda(1+tT^{-1})(1-u_T(t))}{\mu} \right] \quad (7)$$

$$R(\pi, T) = \left\{ \frac{\int_0^t [\lambda(1+tT^{-1})(1-u_T(t))] dt}{\mu} \right\} \quad (8)$$

Solving the equations (2, 3) recursively, using (6, 7, 8), we obtain

$$\begin{aligned} p'_n(\pi) &= -[1 + \rho(\pi, T)] p_n(\pi) + \rho(\pi, T) p_{n-1}(\pi) + p_{n+1}(\pi) \\ p'_0(\pi) &= -\rho(\pi, T) p_0(\pi) + p_1(\pi) \end{aligned} \quad (9-10)$$

Rewritten and Differentiating these equations, we have,

$$\frac{d}{d\pi} Q_0(\pi, T) = -[R(\pi, T)] + \rho(\pi, T) \text{Exp.}\{\pi[1 + R(\pi, T)]\}$$

$$\frac{d}{d\pi} Q_n(\pi, T) = [p'_n(\pi, T) + \{1 + R(\pi, T) + \pi(-[R(\pi, T)] + \rho(\pi, T))\} p_n(\pi)] \text{Exp.}\{\pi[1 + R(\pi, T)]\} \quad (11-12)$$

The initial conditions $[Q_n(t)]_{t=0} = \psi_{jn} : j, n = 0, 1, 2, \dots$ exist and satisfy the system of linear equations. In order to reduce this differential-difference equation (11-12), we introduce the generating function:

$$Q(z, \pi, T) = \sum_{n=0}^{\infty} Q_n(\pi, T) \frac{(z-\pi)^n}{n!} \quad (13)$$

This function will be analytic in z and continuously differentiable in π . Differentiation with respect to z and π gives, using (11-12). Restate the differential difference

equations in term of $\frac{d}{d\pi} Q_n(\pi, T)$, taking summations on both sides with limits $n=0$ to $n=\infty$ as a partial differential equation-w.r.t. π , we get

$$\begin{aligned} \frac{\partial}{\partial \pi} Q(z, \pi, T) &= -\sum_{n=0}^{\infty} Q_n(\pi, T) \frac{(z-\pi)^{n-1}}{(n-1)!} + \sum_{n=0}^{\infty} \frac{(z-\pi)^n}{n!} \frac{\partial}{\partial \pi} Q_n(\pi, T) \\ \frac{\partial}{\partial \pi} Q(z, \pi, T) &= Q_0(\pi, T) + \rho(\pi, T) \sum_{n=0}^{\infty} Q_n(\pi, T) \frac{(z-\pi)^{n+1}}{(n+1)!} \end{aligned} \quad (14-15)$$

Again partial differential equation for $Q(z, \pi, T)$ by using (11-12) w.r.t. z :

$$\frac{\partial^2}{\partial z \partial \pi} Q(z, \pi, T) = \rho(\pi, T) Q(z, \pi, T) \quad (16)$$

These are given by (11-12) and (14-15) which transform in the form of

$$\left[\frac{\partial}{\partial \pi} Q(z, \pi, T) \right]_{z=\pi} = Q_0(\pi, T) \quad (17)$$

$$[Q(z, \pi, T)]_{z=\pi} = Q_0(\pi, T) + \left[\sum_{n=1}^{\infty} Q_n(\pi, T) \frac{(z-\pi)^n}{n!} \right]_{z=\pi} \quad (18)$$

$$Q_j(z, 0, T) = \frac{z^j}{j!} \quad (19)$$

The method of the solution to be used here, we first solve for $Q(z, \pi, T)$ in terms of the function, we define:

$$f_j(\pi, T) = \frac{\partial}{\partial \pi} Q(0, \pi, T) \quad (20)$$

Taking Laplace Transform of (12) and (16) using (20) w.r.t. z and we obtain

$$\frac{\partial}{\partial \pi} Q^*(s, \pi, T) - \frac{\rho(\pi, T) Q^*(s, \pi, T)}{s} = \frac{1}{s} f_j(\pi, T) \quad (21)$$

$$Q^*(s, \pi, T) = \frac{1}{e^{\frac{1}{s}[\pi R(\pi, T)]}} \int_0^{\frac{1}{s}[\vartheta R(\vartheta, T)]} \frac{\pi}{s} f_j(\vartheta, T) e^{-\frac{1}{s}[\vartheta R(\vartheta, T)]} d\vartheta + c \quad (22)$$

$$Q^*(s, \pi, T) = \sum_{n=0}^{\infty} \frac{Q_n(\pi, T)}{n!} \left[c_0^{n(-1)^0} (\pi)^0 \frac{n!}{s^{(n+1)}} + \sum_{k=1}^n (-1)^k c_k^n (\pi)^k \frac{(n-k)!}{s^{(n-k+1)}} \right] \quad (23)$$

At $t=0$, then system is in initial state. Therefore

$$Q^*(s, 0, T) = \sum_{n=0}^{\infty} \frac{Q_n(0, T)}{n!} \frac{n!}{s^{(n+1)}} \quad (24)$$

The solution of (21-23) is

$$Q^*(s, \pi, T) = e^{\frac{1}{s}[\pi R(\pi, T)]} s^{-(j+1)} + \int_0^{\frac{1}{s}[\pi R(\pi, T)]} \frac{\pi}{s} f_j(\vartheta, T) e^{\frac{1}{s}[\pi R(\pi, T)] - \frac{1}{s}[\vartheta R(\vartheta, T)]} d\vartheta \quad (25)$$

Taking Laplace inverse (25) and using the modified Bessel function, with boundary conditions. Then, we get

$$Q_j(z, \pi, T) = [\pi R(\pi, T)]^{-j/2} z^{j/2} I_j[2\{\pi z R(\pi, T)\}^{1/2}] + \int_0^{\pi} I_0[2\{(\pi R(\pi, T) - \vartheta R(\vartheta, T)z)\}^{1/2}] f_j(\vartheta, T) d\vartheta$$

$$Q_j(z, \pi, T) = \Upsilon_j(0, z, \pi, T) + \int_0^{\pi} \Upsilon_0(\vartheta, z, \pi, T) f_j(\vartheta, T) d\vartheta$$

$$\Upsilon_n(\vartheta, z, \pi, T) = [\pi R(\pi, T) - \vartheta R(\vartheta, T)]^{-n/2} z^{n/2} I_n[2\{(\pi R(\pi, T) - \vartheta R(\vartheta, T)z)\}^{1/2}]$$

$$\frac{\partial}{\partial z} \Upsilon_n(\vartheta, z, \pi, T) = \Upsilon_{n-1}(\vartheta, z, \pi, T) \quad (26-28)$$

Noting that

$$\frac{\partial}{\partial \pi} \Upsilon_n(\vartheta, z, \pi, T) = \frac{\partial}{\partial \pi} \left[\frac{[\pi R(\pi, T) - \vartheta R(\vartheta, T)]^{-n/2} z^{n/2} I_n}{[2\{(\pi R(\pi, T) - \vartheta R(\vartheta, T)z)\}^{1/2}]} \right] \quad (29-30)$$

$$\frac{\partial}{\partial \pi} \Upsilon_n(\vartheta, z, \pi, T) = \rho(\pi, T) \Upsilon_{n+1}(\vartheta, z, \pi, T)$$

Define:- $\Upsilon_n(0, 0, 0, T) = \psi_{0n}$

$\Upsilon_{-n}(\pi, \pi, \pi, T) = 0 : n > 0$

$$\Upsilon_0(\pi, \pi, \pi, T) = 1 \quad (31-33)$$

Partial differentiation (27) w.r.t π and using (29-30), (18), we get

$$\left[\frac{\partial}{\partial \pi} Q_j(z, \pi, T) \right]_{z=\pi} = \rho(\pi, T) \Upsilon_{j+1}(0, \pi, \pi, T) + \Upsilon_0(\pi, \pi, \pi, T) f_j(\pi, T) + \int_0^{\pi} \rho(\pi, T) \Upsilon_1(\vartheta, \pi, \pi, T) f_j(\vartheta, T) d\vartheta$$

$$f_j(\pi, T) = Q(\pi, \pi, T) - \rho(\pi, T) \Upsilon_{j+1}(0, \pi, \pi, T) - \int_0^{\pi} \rho(\pi, T) \Upsilon_1(\vartheta, \pi, \pi, T) f_j(\vartheta, T) d\vartheta \quad (34-36)$$

Then,

$$p_n(\pi) = e^{-\pi[1+R(\pi, T)]} \left[\Upsilon_{j-n}(0, \pi, \pi, T) + \int_0^{\pi} \Upsilon_{-n}(\vartheta, \pi, \pi, T) f_j(\vartheta, T) d\vartheta \right] \quad (37)$$

$$f_j(\pi, T) = \omega_j(0, \pi, \pi, T) + \int_0^{\pi} \omega_0(\vartheta, \pi, \pi, T) f_j(\vartheta, T) d\vartheta \quad (38)$$

Here,

$$\omega_j(\vartheta, \pi, \pi, T) = \Upsilon_j(\vartheta, \pi, \pi, T) - \rho(\pi, T) \Upsilon_{j+1}(\vartheta, \pi, \pi, T)$$

Consequently, Equation (37) gives the approximate solution to the single server queue which arrival rate is time dependent and service rate is constant with step function, provided a solution (38) can be found.

Conclusion

In this paper we have obtained the analytic transient solution of M/M/1/N queuing system with a time dependent arrival rate $\lambda(1+tT^{-1})$ using unit step function and service rate μ is constant. This model finds its application in computer-networks, tele-communications, traffic problems and production systems.

References:

1. Abate J. and Whitt W., Transient behavior of the M/M/1 queue, *Queueing Systems*, **2**, 41-65 (1987)
2. Abate J. and Whitt W., Transient behavior of the M/M/1 queue via Laplace transforms, *Advances in Applied Probability*, **20**, 45-178 (1988)
3. Abate J. and Whitt W., Calculating time-dependent performance measures for the M/M/1 queue, *IEEE*, **37**, 1102-1104 (1989)
4. Keller J.B., Time-dependent queues, *SIAM Rev.*, 401-412 (1982)

5. Baily N.T.J., A continuous time treatment of a simple queue using generating function, *J.R.Sata*, **16**, 288-291(1954)
6. Ledermann W. and Reuter G.E.H., Spectral theory for the differential equations of simple birth and death processes, *Philos. Trans. Roy. London*, **246**, 321-369 (1954)
7. Conolly B.W., A difference equation technique applied to the simple queue with arbitrary arrival interval distribution, *J.R.S.S.B*, **21**, 268-275 (1958)
8. Parthasarathy P.R., A transient solution to an M/M/1 queue: a simple approach, *Advances in Applied Probability*,**19**, 997-998 (1987)
9. Clarke A.B., The time dependent waiting line problem, *Uni. of Mic. Engg. Inst.*, Report No: **M720-1,R39** (1953)
10. Koopman B.O., Air terminals queue under time dependent conditions, *Operations Research*, **20**, 1089-1114 (1972)
11. Knessl C., Exact and asymptotic solutions to a PDE that arises in time dependent queues, *Advances in Applied Probability*, **32**, 256-283 (2000)
12. Gross D. and Harris C.M., Fundamentals of queueing theory, *John Wiley, New York*, (2003)
13. Newell G.F., Queues with time-dependent arrival rates-I: the transition through saturation, *Journal Applied Probability*, **5**, 436-451 (1968)
14. Trivedi K.S., Probability and statistics with reliability, queueing and computer science application, *John Wiley*,(2013)
15. Becker L.C., Reducing waiting time at security checkpoints, *Interfaces*, **19**, 57-65 (1999)
16. Saaty T.L., Elements of queueing theory, *McGraw-Hill, New York*, (1961)
17. Collings T., The M/M/ ∞ queue with varying arrival and departure rates, *Operations Research*. **24**, 760-773 (1976)