



Synthesis, Characterization, Antibacterial Activity and DNA cleavage studies of Schiff base Co(II) Transition Metal Complexes

D. Sathis Kumar and Ibrahim Sheriff A.K.*

PG and Research Dept. of Chemistry, C. Abdul Hakeem College (Autonomous), Melvisharam-632509, Tamil Nadu, India
akmhi62@gmail.com

Available online at: www.isca.in, www.isca.me

Received 28th September 2015, revised 4th December 2015, accepted 6th February 2016

Abstract

Three Schiff base ligands L^1 , L^2 and L^3 were achieved by the condensation of *o*-phenylenediamine with *p*-methylbenzaldehyde, *p*-methoxybenzaldehyde and *N*-acetylthiourea respectively. Their complexes with Co(II) were prepared and characterized by elemental analyses, molar conductance, magnetic susceptibility, IR, UV-Vis spectra, NMR, Mass spectral studies, thermal behaviour and Powder X-ray diffraction studies have also been carried out. The IR data determine the bidentate coordination of L^1 , L^2 and tetradentate coordination of L^3 . The XRD data show that Co(II) complexes with L^1 , L^2 and L^3 have the crystallite sizes of 70, 72 and 78 nm respectively. The DNA cleavage activities of the Schiff base and its complexes were monitored by agarose gel electrophoresis method in the presence of hydrogen peroxide and the antibacterial activities of the ligand and complexes have also been carried out.

Keywords: Schiff base complex, Transition metal, Spectral characterization, XRD, DNA cleavage, Antibacterial activity.

Introduction

Schiff bases have been playing an important component in the development of coordination chemistry. Schiff base metal complexes have been studied extensively because of their attractive chemical and physical properties and their wide range of applications in numerous scientific areas. They play an important role in both synthetic and structural research, because of their preparative accessibility and structural diversity¹. Metal complexes of Schiff base derived from the reaction of *o*-phenylenediamine with aliphatic and aromatic amines represent a series of compounds containing nitrogen, sulphur and oxygen donor atoms that has been widely studied. Schiff base molecules afford potential sites for bio chemically active compounds that are related to intermolecular hydrogen bonding and proton transfer equilibria². Schiff bases of *o*-phenylenediamine and its complexes have a variety of applications including biological, analytical and clinical^{3,4}.

The Schiff base ligand is found to be dibasic with N_2 and N_2S_2 donor sites. The ¹H-NMR, ¹³C-NMR and Mass spectra confirm the formation of N_2 and N_2S_2 bidentate and quadridentate Schiff base chelate ligands. All the metal complexes have been fully characterized with the help of elemental analysis, molar conductance, magnetic susceptibility and spectral studies. The analytical data helped to elucidate the structure of the metal complexes. The complexes exhibit electrolytic behavior in ethanol. The IR spectral data suggest the coordination of azomethine with the central metal ion. The magnetic data suggest an octahedral geometry for Co(II) metal complexes. The thermogravimetric analysis shows that Co(II) metal complexes possess two aqua molecules coordinated in an octahedral

environment. The complexes were subjected to powder XRD patterns with PAN-analytical diffractometer with Cu-K α radiation of wavelength 1.54060 Å operating at a voltage of 30 kV and a current of 40 mA. The nucleolytic cleavage activities of the complexes were assayed on pUC19 DNA using gel electrophoresis in the presence of H₂O₂ and the complexes show promising nuclease activity. The compounds were subjected to antimicrobial activity screening and Minimum Inhibitory Concentration (MIC) is determined. The complexes show significant growth inhibitory activity against the bacteria like *Bacillus cereus*, *Streptococcus pyogenes*, *Listeria monocytogens*, *Escherichia coli* and *Salmonella typhi* than the free ligands. The MIC of 31.25 µg/ml of complex of L^3 is noticeable.

Materials and Methods

Experimental: All the reagents were of AR grade and the solvents were purified by standard methods. IR spectra (4000 – 400cm⁻¹) taken on KBr disc using a Perkin Elmer Spectrum ONE- N017-1159 Spectrophotometer. Micro analysis of carbon, hydrogen, nitrogen and sulphur were obtained using elemental analyzer, magnetic moments were measured at room temperature on Vibrating Sample Magneto meter EG and G Model: 155 using Hg [Co(CN)₄] as standard. Electrical conductances of the complexes were made on a systronic conductivity meter type 304 in DMSO with a dip type cell having platinum electrode. The UV- visible spectra were run on a Hitachi U-2800 spectrophotometer (200-1100 nm) in nujol mull. TGA-DTA analyses were obtained by using NETZCH STA-409C/CD thermal analyzer. The ¹H NMR, ¹³C NMR spectra were recorded on JKM-ECS 400 in DMSO-d₆ solvent.

The mass spectrum was recorded using WATERS-Q-T of premier- HAB213, electro spray ionization-MS.


Antibacterial activity and MIC values: The antibacterial screening of the compounds was carried out against the Gram positive bacteria (*Bacillus cereus*, *Streptococcus pyogenes* and *Listeria monocytogens*) and Gram negative bacteria (*Escherichia coli* and *Salmonella typhi*) on the free ligand and metal complexes by cup plate method using the nutrient agar as medium. In a typical procedure, molten nutrient agar kept at 115°C –120°C was poured into a Petri dish and allowed to solidify. The small wells (10 mm diameter with 1cm distance) were made in the agar medium by carefully using a sterile cork and these were completely filled with test solutions. The plates were incubated for 24 hours at 37 °C. The diameters of the zones of inhibition for all the test compounds were measured and the results were compared with that of standard *Tetracycline* at the same condition. The antibacterial activities were done at 15.62, 31.25, 62.5 and 125 µg/ml concentration in DMSO solvent by using three G +ve and two G –ve bacteria by the MIC method.

DNA cleavage studies: The pUC19 DNA at pH 7.5 in Tris-HCL buffered solution was used to form agarose electrophoresis. Oxidative cleavage of DNA was examined by keeping the concentration of the 30 µM of complex and 2 µL of pUC19 DNA and made up the volume to 16 µL with 5mM Tris-HCL/ 5mM NaCl buffer solution. The resulting solution was incubated at 37°C for 2 h and electrophoresed for 2 h at 50V in Tris-acetate-EDTA (TAE) buffer using 1% agarose gel containing 1.0 µg/ml ethidiumbromide and photographed under UV light. All the experiments were executed at room temperature.

Synthesis of Schiff base L¹, L², L³: A solution of o-phenylenediamine (0.01mole) and p-methylbenzaldehyde / p-methoxy benzaldehyde / N-acetylthiourea (0.02mole) in 1:2 molar ratio in ethanol was mixed with constant stirring. The resulting mixture was refluxed for five hours in a water bath. The concentrated solution was filtered out, washed with ethanol and recrystallized using a solution of chloroform and ethanol mixture, its purity was checked by TLC.

Synthesis of metal complexes (ML¹, ML², ML³): A hot ethanolic solution of Schiff base (L¹/L²) with 1, 1'-binaphthyl-2, 2'-diol (BINOL) and an ethanolic solution of metal acetate solution were mixed in 1:1:1 molar ratio. The mixture was refluxed for about 2-4 h in water bath. The mixture was cooled in ice. The resulting precipitate was then filtered, washed with ethanol and recrystallized using a solution of chloroform and ethanol mixture, its purity was checked by TLC. The above procedure followed by ML³ complex was prepared by (L³) ligand and Co(II) in 1:1 molar ratio^{5,6}.

o-phenylenediamine + p-methylbenzaldehyde
o-phenylenediamine + p-methoxy benzaldehyde
o-phenylenediamine + N-acetylthiourea



L¹ (Figure-1)
L² (Figure-2)
L³ (Figure-3)

Results and Discussion

Micro analysis: Co(II) complexes are stable at room temperature, insoluble in water but soluble in DMF and DMSO. The physical properties and analytical data of the ligands and their complexes are given in Table-1. Elemental analysis data of the complexes are in good agreement with theoretical values. The analytical data indicate the molecular formula of the complexes as [Co L¹ (BINOL) (H₂O)₂], [Co L² (BINOL) (H₂O)₂] and [Co L³ (H₂O)₂]. The Co(II) complexes have lower molar conductance values indicating that the L¹ and L² complexes are non-electrolytes⁷. The L³ Co(II) complexes have higher molar conductance value which shows that it acts as in 1:2 electrolyte⁸.




Figure-1
Schiff base free ligand (L¹)




Figure-2
Schiff base free ligand (L²)




Figure-3
Schiff base free ligand (L³)

Table-1
Physical and analytical data of Schiff bases and their metal complexes

Ligand /complex	M.F (M.Wt.)	Yield %	Found (Calcd.) %					Λ_m Scm ² Mol ⁻¹
			C	H	N	S	Co	
C ₂₂ H ₂₀ N ₂ (L ¹)	312.42	72	84.58 (84.52)	6.45 (6.40)	8.97 (8.94)	-	-	-
C ₂₂ H ₂₀ N ₂ O ₂ (L ²)	344.41	75	76.72 (76.70)	5.85 (5.82)	8.13 (8.10)	-	-	-
C ₁₂ H ₁₆ N ₆ S ₂ (L ³)	308.43	70	46.73 (46.70)	5.23 (5.19)	27.25 (27.20)	20.79 (20.74)	-	-
[Co L ¹ (BINOL) (H ₂ O) ₂]	691.69	76	72.93 (72.90)	5.25 (5.21)	4.05 (4.00)	-	8.52 (8.48)	44.10
[Co L ² (BINOL) (H ₂ O) ₂]	723.69	80	69.71 (69.67)	5.02 (4.98)	3.97 (3.92)	-	8.14 (8.12)	46.30
[Co L ³ (H ₂ O) ₂]	521.48	77	36.85 (36.81)	5.03 (4.97)	16.12 (16.08)	12.30 (12.27)	11.30 (11.28)	146.05

Electronic Spectra: The electronic absorption spectra of free Schiff base ligands (L¹, L² and L³) and their Co(II) complexes were recorded in ethanol solution (10⁻³ M) in the range 200–1000 nm at room temperature Table-2. Schiff base ligands show two bands at 241–284 nm and 305–378 nm corresponding to $\pi \rightarrow \pi^*$ transition of aromatic benzene moiety and $n \rightarrow \pi^*$ transition of azomethine C=N bond respectively.

All the complexes exhibit three bands at 10,690 – 13,100 cm⁻¹ (ν_1), 15,185 – 16450 cm⁻¹ (ν_2) and 19234 – 21760 cm⁻¹ (ν_3) which correspond to ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$, ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$ and ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$ respectively. The observed magnetic moment values 4.66 BM, 4.56 BM and 4.58 BM indicate that the Co(II) complexes are paramagnetic having of three unpaired electrons in an octahedral environment^{9,10}.

Infrared Spectra: The Schiff base ligands L¹, L² and L³ show $\nu(C=N)$ azomethine bands observed at 1643, 1619 and 1689 cm⁻¹. On complexation, this band was shifted to 1617, 1611 and 1638 cm⁻¹ regions due to the coordination of azomethine nitrogen^{11, 12} to the Co(II) ion Table-3. In the Schiff base ligand (L³) the N-acetylthiourea $\nu(C=S)$ band appearing at 1042 cm⁻¹. The band was shifted to lower frequency (1018 cm⁻¹) on complexation indicating that the sulphur is coordinated to the Co(II) ion. The new band appears at 410 cm⁻¹ due to coordination of M-S. Furthermore, the presence of coordinated water molecules is confirmed by the bands at 3450, 3408 and 3412 cm⁻¹ in CoL¹, CoL² and CoL³ complexes may respectively attributed to O-H stretching vibrations. In all these complexes two new bands appear at in the range 446 – 492 cm⁻¹ and 526 – 545 cm⁻¹ respectively the existence of $\nu(M-N)$ and $\nu(M-O)$ bonds.

¹H-NMR spectrum of Schiff base: The signals observed in the ¹H-NMR spectra of the Schiff bases under study Figure-4. ¹H-

NMR spectral data of the Schiff base (L¹) was recorded in CDCl₃, shows peaks at 7.58-6.97 and 2.38-2.32 ppm due to aromatic protons and CH₃ group respectively Table-4. The formation of Schiff base (L²) was further confirmed by the ¹H-NMR spectra. ¹H-NMR spectra of the ligand were taken in DMSO-d₆ solvent Figure-5. The aromatic region gives a set of multiples in the range 6.8–7.6 ppm for the Schiff base ligand, while the azomethine protons were observed in the range 8.4–8.6 ppm. The methoxy group protons show a peak at 3.6-3.8 ppm. The formation of Schiff base (L³) ligand was further confirmed by the ¹H-NMR spectra. ¹H-NMR spectra of the ligand were taken in DMSO-d₆ solvent Figure-6. The aromatic region gives a set of multiples in the range 7.3-7.6 ppm. The methyl group protons show a peak at 2.09 ppm. NH group protons show a peak at 2.06 ppm. NH₂ group protons show a peak at 9.72 ppm. The total number of protons present in the Schiff base exhibited signals their respective in their expected region. It was also observed that DMSO did not show any coordinating effect on all the ligands¹³.




Figure-4
¹H-NMR spectrum of the free Schiff base (L¹)

Table-2
UV-Vis spectra of Schiff base and its complexes

Ligand / Complex	Absorbance nm	ν / cm^{-1}	Assignment	Geometry	Magnetic moment (BM)
L^1	241	41,150	$\pi \rightarrow \pi^*$	-	-
	378	26,455	$n \rightarrow \pi^*$		
L^2	243	41,152	$\pi \rightarrow \pi^*$	-	-
	305	32,786	$n \rightarrow \pi^*$		
L^3	284	35,211	$\pi \rightarrow \pi^*$	-	-
	337	29,673	$n \rightarrow \pi^*$		
[Co L^1 (BINOL) (H ₂ O) ₂]	790	12,650	${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$	Octahedral	4.66
	626	15,960	${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$		
	460	21,760	${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$		
[Co L^2 (BINOL) (H ₂ O) ₂]	763	13,100	${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$	Octahedral	4.56
	608	16,450	${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$		
	467	21,420	${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$		
[Co L^3 (H ₂ O) ₂]	935	10,690	${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$	Octahedral	4.58
	659	15,185	${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$		
	520	19,231	${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$		

Table-3
IR Spectral data of ligands and its metal complexes (cm^{-1})

Ligand / Complex	$\nu_{\text{O-H}_2}$	$\nu_{\text{C=N}}$	$\nu_{\text{C=S}}$	$\nu_{\text{M-S}}$	$\nu_{\text{M-N}}$	$\nu_{\text{M-O}}$
L^1	-	1643.28	-	-	-	-
L^2	-	1619.52	-	-	-	-
L^3	-	1689.52	1042.45	-	-	-
[Co L^1 (BINOL) (H ₂ O) ₂]	3450.14	1617.88	-	-	545.12	492.22
[Co L^2 (BINOL) (H ₂ O) ₂]	3408.48	1611.31	-	-	536.75	461.38
[Co L^3 (H ₂ O) ₂]	3412.02	1638.54	1018.08	410.08	526.26	446.34

Table-4
 ${}^1\text{H-NMR}$ Spectral data of ligand and its metal complexes (δ ppm) in d_6 -DMSO

Ligand	${}^1\text{H-NMR}$ Spectra	${}^{13}\text{C-NMR}$ spectra
L^1	8.32 (2H,s, CH=N), 6.97-7.58(12H,m, Ar-H), 2.32-2.38(6H,-CH ₃)	21.1 -21.1(2C,-CH ₃), 121.8,124.6,130.2,134.5,141.3 (18C, Ar-C), 161.5 (2C, CH=N)
L^2	8.60 (2H,s,CH=N), 6.8-7.6(12H,m, Ar-H), 3.80(6H,-OCH ₃)	54.68-55.60(2C,-OCH ₃), 114.7,121.5,123.8,127.8,129.6,131.4,162.8 (18C, Ar-C), 161.8(2C, CH=N)
L^3	2.09-2.15(6H,s, -CH ₃), 2.06-2.14(2H, -NH), 9.72(4H, -NH ₂), 7.29-7.65(4H, Ar-H)	16.7(2C, -CH ₃), 123.5, 131.2, 141.6 (6C, Ar-C), 163.8 (2C, C=N), 184.7(2C, -C=S)

Table-5
Determination of MIC for antibacterial activity

Micro organisms	L ¹				M ¹ L ¹				L ²				M ² L ²				L ³				M ³ L ³			
	a	b	c	d	a	b	c	d	a	b	c	d	a	b	c	d	a	b	c	d	a	b	c	d
<i>Bacillus cereus</i>	-	+	+	+	-	-	+	+	-	+	+	+	-	+	+	+	-	+	+	+	-	-	+	+
<i>Streptococcus pyogenes</i>	-	-	+	+	-	-	+	+	-	+	+	+	-	+	+	+	-	-	-	+	-	-	+	+
<i>Listeria monocytogens</i>	-	-	+	+	-	+	+	+	-	+	+	+	-	+	+	+	-	-	+	+	-	-	+	+
<i>Escherichia coli</i>	-	-	+	+	-	-	+	+	-	-	+	+	-	-	+	+	-	+	+	+	-	+	+	+
<i>Salmonella typhi</i>	-	-	+	+	-	+	+	+	-	-	+	+	-	-	+	+	-	-	+	+	-	+	+	+

a= 125 µg/ml, b= 62.5 µg/ml, c= 31.25 µg/ml, d= 15.62 µg/ml: M= Co(II) ion, Minus (-) indicates the absence of growth, Plus (+) indicates presence of growth

Table-6
Antibacterial activities MIC values

Micro organisms	MIC values (µg/ml)					
	L ¹	M ¹ L ¹	L ²	M ² L ²	L ³	M ³ L ³
<i>Bacillus cereus</i>	125	62.5	125	125	125	62.5
<i>Streptococcus pyogenes</i>	62.5	62.5	125	125	31.25	62.5
<i>Listeria monocytogens</i>	62.5	125	125	125	62.5	62.5
<i>Escherichia coli</i>	62.5	62.5	62.5	62.5	62.5	125
<i>Salmonella typhi</i>	62.5	125	62.5	62.5	62.5	125




Figure-5

¹H-NMR spectrum of the free Schiff base (L²)




Figure-6

¹H-NMR spectrum of the free Schiff base (L³)

¹³C-NMR spectrum of Schiff base: The signals observed in the ¹H -NMR spectra of the Schiff bases under study Figure-7. ¹³C- NMR signals of the Schiff base (L¹) was recorded in CDCl₃ shows peaks at 121.8 – 141.3 and 21.1 -21.5 ppm due to aromatic carbons and CH₃ groups respectively. The formation of Schiff base ligand (L²) was confirmed by the ¹³C-NMR spectra. ¹³C-NMR spectra of the ligand were taken in DMSO-d₆

solvent Figure-8. The aromatic region shows a set of multiples in the range 114.13-162.8 ppm for the Schiff base ligand, while the azomethine carbons were observed in the range 161.8 ppm. The methoxy group carbon shows appear in the range at 54.6 – 55.6 ppm. The formation of Schiff base ligand (L³) was confirmed by the ¹³C-NMR spectra Figure-9. The aromatic region gives a set of multiples in the range 123.5 - 141.6 ppm for the Schiff base ligand, while the azomethine carbons were

observed in the range 163.8 ppm. The methyl group carbons appear at 16.7 ppm. The C=S group carbon show a peak at 184.7ppm. All the total number of carbons present in the Schiff base exhibited signals their expected regions¹⁴.




Figure-7
¹³C-NMR spectrum of the free Schiff base (L¹)




Figure-8
¹³C-NMR spectrum of the free Schiff base (L²)

Mass spectral fragmentation of the Schiff base: The mass spectrum of Schiff base (L¹) shows a base peak m/e+ at 313 which is due to the observed molecular weight of the prepared Schiff base Figure-10. This proves the condensation of aldehyde and amine to form the Schiff base^{15,16}. The mass spectra of the ligand recorded at room temperature were used to ascertain the stoichiometric composition. The Ligand (L²) shows a molecular ion peak at m/z 345, which corresponds to [L+H]⁺ peak as the calculated m/z being 344 Figure-11. The Ligand (L³) shows a molecular ion peak at m/z 309, which corresponds to [L+H]⁺ peak as the calculated m/z being 308 Figure-12.

Thermal studies: The TGA curve of the cobalt complex [Co L¹ (BINOL) (H₂O)₂] shows that it is stable up to 115°C. A weight loss is observed around 128-135°C corresponding to the elimination of two coordinated water molecules^{17, 18}. In the complexes [Co L² (BINOL) (H₂O)₂] and [Co L³ (H₂O)₂] this loss of coordinated water is observed at 125-140 and 115-130°C




Figure-9
¹²C-NMR spectrum of the free Schiff base (L²)

respectively.

Powder X-ray diffraction studies: The XRD pattern of Co(II) complexes show well defined crystalline peaks indicating that the samples are crystalline in nature Figure-13. The above complexes have specific 'd' values which can be used for its characterization. The crystallite size of the complexes d_{XRD} could be estimated from XRD patterns by the Scherrer's formula $d_{XRD} = 0.9\lambda/\beta (\cos\theta)$, where λ is the wavelength, β is the full width at half maxima and θ is the diffraction angle. The XRD shows that Co(II) complexes have the average crystallite size of 70, 72 and 78 nm respectively, suggesting the complexes to be nanocrystalline^{19,20}.

DNA cleavage studies: Gel electrophoresis experiments were performed using pUC19 DNA with ligands and their complexes in the presence and absence of H₂O₂. Complexes exhibit cleavage capability at low concentration (40 μM). The ligand exhibits no significant activity in the presence of oxidant. The activity was much higher for the complexes in presence of H₂O₂. When the calf-thymus DNA is subjected to electrophoresis, relatively fast migration will be observed for the intact super coil form (Form I). If scission occurs on one strand (nicking), the super coil will relax to generate a slower moving open circular form (Form II). If both strands are cleaved, a linear form (Form III) that migrates between Forms I and II will be generated^{21,22}. The complexes show more activity in the presence of oxidant which may be due to the reaction of hydroxyl radical with DNA. These hydroxyl free radicals participate in the oxidation of the deoxyribose moiety followed by hydroxyl cleavage of sugar phosphate backbone. The results of DNA cleavage studies have been shown in Figure-14. All metal complexes were able to convert DNA (Form I) into open circular (Form II). The L¹ Co(II) complex was found to be highly active in cleaving DNA in the presence of hydrogen peroxide. The H₂O₂ is coordinated to the cobalt ion of the complex, affording a peroxo - cobalt species. This coordinated peroxide ion attacks the DNA phosphate bond via a nucleophilic mechanism and hydrolyzes the P-O bond²³.




Figure-10
Mass spectra of the Schiff base (L¹)




Figure-12
Mass spectra of the Schiff base (L³)




Figure-11
Mass spectra of the Schiff base (L²)




Figure-13
XRD pattern of Co(II) complexes




Figure-14
DNA cleavage Schiff base metal complexes





Figure-15
Antibacterial activity



Antibacterial activity: The antibacterial activity of the Co(II) complexes were studied against five pathogenic bacteria strains. The results of the antibacterial activity studies for the complexes and standard antibiotic *tetracycline* were evaluated by Muller Hinton agar diffusion method and serial dilution sensitivity test against both Gram-positive and Gram-negative bacteria. DMSO solvent was also used as control. As it could be seen from these tables, both Gram positive and Gram negative bacteria were affected by these antibacterial agents and *L.monocytogenes* was the most sensitive microorganism of the studied complexes Figure-15. On the other hand, *E.coli*, *S. typhi*, *B.cereus*, and *S.pyogenes* were, to some extent, more resistant to these compounds. It is previously reported that the presence of electronegative and electron donating groups on the sulphur part

of the Schiff base increases the antibacterial activity of their metal complexes^{24,25}.

The MIC was calculated as the highest dilution showing complete inhibition of the tested strains and reported in Table-5 and 6. These Schiff base and complexes also disturb the respiration process of the cell and thus block the synthesis of proteins, which further restricts the growth of the microorganisms. The ML³ complex has good activity when compared to Schiff base and other metal complexes.²⁶

Conclusion

In this paper, we presented three Schiff base ligands from o-

phenylenediamine and p-methylbenzaldehyde/p-methoxy benzaldehyde / N-acetylthiourea and their Co(II) complexes have been synthesized using the Schiff base ligands. The ligands and complex were characterized by spectral and analytical data. The electronic spectral data indicates that all the Co(II) complexes are paramagnetic with octahedral geometry. The pUC19 DNA cleavage studies explain the total cleavage of pUC19 DNA by Co(II) complexes in the presence of H₂O₂. The antibacterial studies fulfill with the complexes confirm that they are good antibacterial agents. Their MIC values being 125 µg / litre.

Acknowledgements

The authors express their sincere thanks to the management, PG and Research department of Chemistry C. Abdul Hakeem College, Melvisharam. We also thank SAIF, IIT Kanpur for providing FTIR, TGA-DTA analyzer, Mass, SAIF, IIT Chennai for magnetic susceptibility data, VIT, Vellore for providing Electronic spectral studies, DNA cleavage, antibacterial activity and powder X-ray diffraction data.

References

1. Selwin Joseyphus R and Sivasankaran Nair M. (2010). Synthesis, characterization and biological studies of some Co(II), Ni(II) and Cu(II) complexes derived from indole-3-carboxaldehyde and glycylglycine as Schiff base ligand, *Arabian Journal of Chemistry*, 3, 195–204.
2. Prabakararishnan R, Manoranjitham S and Geetha Kannappan. (2014). Synthesis of Copper (II) Complexes Using Pentadentate Schiff Base Ligand By Eco Friendly Solventless Method and Its Antimicrobial Activities, *World journal of pharmaceutical research.*, 3(9), 1286-1299.
3. Nagajothi A, Kiruthika A, Chitra S and Parameswari K. (2012). Synthesis and Characterization of Tetradentate Co(II) Schiff Base Complexes : Antimicrobial and DNA Cleavage Studies, *International Journal of Research in Pharmaceutical and Biomedical Sciences.*, 3(4), 1768-1778.
4. Sreenivas V, Srikanth G, Aruna M, Vijaya Kumar P, Muralidhar Reddy P and Ravinder V. (2014). Synthesis, Characterization and Antibacterial Activity and DNA cleavage Studies of tetra dentate Schiff bases and their Zn (II) Complexes, *Research Journal of Chemical Sciences*, 4(6), 66-72.
5. Peiyuan Wang, Jie Yang, Jian Liu, Lei Zhang and Qihua Yang. (2009). Chiral mesoporous organosilicas with R-(+)-Binol integrated in the framework, *Microporous and Mesoporous Materials.*, 117, 91–97.
6. Barbara Paul, Christian Na`ther, Katharina M. Fromm and Christoph Janiak. (2005). Chiral S-1,1'-bi-2-naphthol (S-BINOL) as a synthon for supramolecular hydrogen-bonded {(S-BINOLATⁿ)(S-BINOL)_n} strands with naphthyl-paneled cavities or channels for a Cd(NH₃)₄-fragment (n =2) or [Ag(NH₃)₂]⁺ (n =1). Part 2⁺, *Cryst Eng Comm.*, 7(51), 309–319.
7. Sathis Kumar D., Rajiv Gandhi S. and Ibrahim Sheriff A.K. (2015). Synthesis, spectral characterization and antimicrobial activity of bidentate Schiff's base (N₂) transition metal complexes, *Journal of Chemical and Pharmaceutical Research*, 7(1), 416-423.
8. Imran Ali A, Waseem A., Wani A and Kishwar Saleem. (2013). Empirical Formulae to Molecular Structures of Metal Complexes by Molar Conductance, *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, 43, 1162–1170.
9. Vogel AI A, Textbook of Quantitative Inorganic Analysis Including Elementary Instrumental Analysis, fourth ed. Longman, London (1978)
10. Lever A.B.P. (1984). Inorganic electronic spectroscopy, 2nd edition Elsevier, New York.
11. Sathis kumar D., Gnanavel S., Rajiv Gandhi S., Anbazhagan Mageswari, Sivashanmugam Karthikeyan and Ibrahim Sheriff A.K. (2015), Synthesis, Spectral Characterisation and Antibacterial activity of Some Novel Schiff Base Chelate (N₂) Complexes of Transition Metal Ions, *International Journal of Frontiers in Science and Technology.*, 3(1), 72-81.
12. Nakamoto K. (1997). Infrared and Raman Spectra of inorganic and coordination compounds, 5th Edn, Part A and B. Wiley, New York.
13. Prasad A.V.G.S, Venkateswara Rao I P and Prasad P.S.S. (2014). Synthesis, Characterization, Antifungal, Antibacterial Studies of Heterocyclic Schiff Base Metal Complexes, *International Journal for Pharmaceutical Research Scholars*, 3(1), 63-68.
14. Ali MA, Mirza AH and Butcher RJ. (2001). Synthesis and characterization of copper(II) complexes of the methylpyruvate Schiff base of S-methyldithiocarbamate (Hmpsme) and the Xcrystal structures of Hmpsme and [Cu(mpsme)Cl], *Polyhedron.*, 20, 1037–1043.
15. Babu K and Pitchai P. (2013). Synthesis, characterization biological studies of new bis mannich base and its transition metal complexes, *International Journal of Chemical and Pharmaceutical Sciences*, 2013 4(4), 39-44.
16. Singh DP and Rana VB. (1995). Binuclear chromium (III), manganese (III), iron (III) and cobalt (III) complexes bridged by diaminopyridine, 14, 2901-2906.
17. Saydam S and Alakn C. (2001). Polish, Synthesis and characterization of a new thiazole Schiff base 5-bromo-2-hydroxybenzylidene-2-aminobenzothiazole and its complexes with Co(II), Cu(II) and Ni(II), *J Chem.*, 75,

- 29–33.
18. Enis Nadia Md Yusof, Thahira Begum S.A. Ravoof, Edward R T. Tiekink, Abhimanyu Veerakumarasivam, Karen Anne Crouse, Mohamed Ibrahim Mohamed Tahir and Haslina Ahmad. (2015). Synthesis, Characterization and Biological Evaluation of Transition Metal Complexes Derived from N, S Bidentate Ligands, *Int. J. Mol. Sci.*, 16, 11034-11054.
 19. Sukdolak S, Vukovi N, Soluji S, Miloev M, Manojlovi N and Krsti J. (2009). Synthesis, Spectroscopic Investigation and Antiactivity Activity of Schiff Base Complexes of Cobalt (II) and Copper (II) Ions, *Rasayan journal chemistry.*, 2(2), 261-266.
 20. Geary WJ. (1971). The use of conductivity measurements in organic solvents for the characterisation of coordination compounds, *Coord. Chem. Rev.*, 7(1), 81-122.
 21. Dhanaraj C.J. and Nair M.S. (2009). Synthesis and characterization of metal(II) complexes of poly(3-nitrobenzylidene-1-naphthylamine-co-succinic anhydride), *Eur. Polym. J.*, 45(2), 565–572.
 22. Kannan D and Arumugham MN (2012). Synthesis, Characterisation, DNA-Binding Studies and antimicrobial activity of Copper(II) Complex with 1,10 Phenanthroline, L-Tyrosine and Thiourea as Ligands, *International Journal of Research in Controlled Release.*, 2(4), 10-17.
 23. Raman N, Dhaveethu Raja J and Sakthivel A (2007). Synthesis, spectral characterization of Schiff base transition metal complexes: DNA cleavage and antimicrobial activity studies, *J Chem Sci.*, 119, 303-310.
 24. Sitlani AS, Long EC, Pyle AM, Barton JK, DNA photocleavage by phenanthrenequinone diimine complexes of rhodium (III): shape-selective recognition and reaction, *J. Am. Chem. Soc.*, 114, 2303–2312 (1992)
 25. Kalia SB, Lumba K, Kaushal G and Sharma M. (2007). Magnetic and spectral studies on Co(II) chelates of a dithiocarbazate derived from isoniazid, *Ind J chem.*, 46A, 1233-1239.
 26. Braide W, Nwaoguikpe RN, Oranusi SE, Udegbonam LI, Akobondu C and Okorundu SI (2011). The effect of biodeterioration on the nutritional composition and microbiology of an edible long-winged reproductive termite, *Macrotermes bellicosus*. Smeathman, *Internet Journal of Food Safety*, 13, 107-114.