



Influence of abattoir wastes on soil microbial and physicochemical properties

Emmanuel Dan^{1*}, Kayode Fatunla² and Solomon Shuaibu¹

¹Department of Chemistry, University of Uyo, Uyo, Akwa Ibom State, Nigeria

²Microbiology Department, University of Uyo, Uyo Akwa Ibom State, Nigeria
danmarble2003@gmail.com

Available online at: www.isca.in, www.isca.me

Received 22nd August 2018, revised 20th November 2018, accepted 15th December 2018

Abstract

*Influence of abattoir wastes on physicochemical and microbiological properties of soil samples obtained from Akwa Ibom State, Southern Nigeria were investigated using standard analytical and microbiological methods. Studied abattoirs and the control soils were in the sandy-clay-loamy soil category with varied quantities of sand, silt and clay. Bulk density, pH, electrical conductivities, salinity, moisture content, total organic and carbon content, cation exchange capacities, total petroleum hydrocarbon, nitrogen and phosphorus contents of studied abattoir soils were higher than in control. Essential elements (K, Na, Ca, and Mg) and trace metals (Fe, Zn, Cd, Cu, Pb, Cr and Ni) levels were also higher in abattoir soils than in control though were within permissible limit in soil except for Fe. Metal pollution index (MPI), enrichment factor (EF), geo-accumulation index (Igeo), degree of contamination (Cdeg) and pollution load index (PLI) of trace metals have also been calculated using existing pollution models. Microbial studies revealed total heterotrophic bacteria ranged from 6.41 ± 0.43 to $7.91 \pm 0.58 \log_{10} \text{CFU/g}$ while fungal count ranged from 4.94 ± 0.26 to $5.79 \pm 0.34 \log_{10} \text{CFU/g}$. Among the four (4) locations, IK2 had the highest heterotrophic bacterial densities of $7.91 \pm 0.58 \log_{10} \text{CFU/g}$ while IK1 had the highest fungal count of $5.79 \pm 0.34 \log_{10} \text{CFU/g}$. A total of six (6) bacteria (*Klebsiella*, *Micrococcus*, *Pseudomonas*, *Bacillus*, *Escherichia* and *Enterobacter*) and two (2) fungi (*Aspergillus* and *Penicillium*) species were isolated. The study revealed a significant ($p=0.05$) increase in the number and varieties of microorganisms most of which may be pathogenic but are more often than not indicators of recent faecal pollution in the soil impacted with the abattoir wastes.*

Keywords: Microbial, physicochemical properties, abattoir waste, pathogens.

Introduction

Studies have shown that environmental pollution and its attendant problems on land, air and water qualities are severe now than before. There are several evidences to this fact ranging from soil fertility loss, depletion of biodiversity, several health problems (those leading to metabolic disorder), ecological effect and others¹⁻³.

The major cause of these pollutions is the indiscriminate discharge of wastes into these natural habitats thereby tampering with the natural workings of the environment. Solid and liquid wastes are usually disposed off on open landfills, waterways, rivers and stream indiscriminately by most industries (small and large), and the entire populace. These practices are very common in Nigeria till date since there are no well define management protocols on solid waste disposal.

Abattoir waste is another emerging solid waste whose rate of generation is becoming alarming. Abattoir or slaughter house is a dedicated place usually outside a residential area where meat processing is usually carried out. According to Hornby⁴; it is also a place for killing and processing of meat. Several activities are involved in processing of meat from receiving and holding of livestock to final packaging (ready to cook)⁵. However, in Nigeria, abattoir activities are operated by butchers who have

little or no idea of sanitary principles. These activities are usually accompanied by the generation of large amounts of organic and inorganic wastes which are hitherto discharged into soils and water bodies around the abattoir premises^{6,7}.

It is not possible to know the exact composition of abattoir wastes since it contain so many components, and is usually detrimental to any environment where they are discharged. Various organs of cattle like muscles, blood, liver and kidney have been reported to contain trace metals, faeces of livestock consist of mucus, bacteria, cellulose fibre, paunch manure which is very acidic in nature and others⁸⁻¹¹. Additional reports have been made on the effect of abattoir wastes on soil including increase concentration of trace metals, increase population of decomposers, loss of aesthetic value, excessive soil nutrient enrichment and increase toxin accumulation, as well as large accumulation of sulphides, mercaptans, amines and organic acids^{3,11-15}.

In Akwa Ibom State, most abattoirs have farmlands surrounding them and these farmlands are usually cultivated by local farmers living in the vicinity of these abattoirs. These soils are used to plant crops mostly vegetables and legumes which are consumed by humans due to good crop yields from these abattoir soils without any assessment of its sanitary nature. Therefore this study seeks to investigate the influence of abattoir wastes on the

physicochemical properties, trace metal levels and diversity of microorganisms in such soils.

Materials and methods

Sample collection and pre-treatment: Abattoir soil samples (0–20cm) were collected from four (4) different abattoir soil locations, two each from Ikot Ekpene (IK1, IK2), and Eket Local Government Area (EK1, EK2) respectively in Akwa Ibom State, Southern Nigeria using Soil Auger. At each Local Government Area, soil samples of the same depth were also collected from soils not close to a slaughter house or abattoir waste site and used as Control samples (IKC, EKC). The abattoir soils were collected after clearing off waste materials from the soil within the vicinity of these abattoirs and were done in January 2018. A total of six (6) soil samples were collected for this study. Soil samples for microbial studies were put in a polythene bags already sterilized and taken to the microbiology laboratory for immediate analysis.

Determination of essential and trace elements: Abattoir and Control soils were first air dried for three (3) days to drive off any liquids, before being ground using mortar and pestle to achieve homogeneity. Prior to atomization for the total trace elements (Fe, Zn, Cd, Cu, Pb, Cr, Mg, and Ni) determination, 1.00g sample each was placed in a crucible and 10ml Aqua Regia (HCl/HNO₃ 3:1) was added. Resulting solutions were then left to digest under reflux for an hour and then evaporated to dryness at a temperature of 120°C on a hot plate under a fume hood. The samples were then leached with 5ml HCl and made up to a 10 ml mark before transferring them quantitatively into small sample bottles for atomization using inductively couple plasma optical emission spectrophotometer (Agilent 710). Na, K and Ca were determined using flame emission spectrophotometer (Model 381 and 391) at different wavelengths.

Determination of textural class and physicochemical properties: Particle size distribution through the physical analytical test was carried out using the hydrometer methods¹⁶. Textures class of the abattoir soils and Control were determined using United States Department of Agriculture (USDA) textural triangle, while Bulk densities were carried out using gravimetric method as described by Omofunmi *et al*¹⁷. The pH and electrical conductivity of studied abattoir soils and Control were measured using portable meters after calibration with standard solutions¹⁸. Salinity was determined following the method of Edori *et al*¹⁹. Moisture content was determined gravimetrically after drying the soils in an oven (Gallenkamp OV-330) at 105°C until a constant weight was obtained. Total organic matter contents were measured by Walkely-Black's wet oxidation method described in details by Ebong *et al*²⁰. Total organic carbon and cation exchange capacity were determined following the methods of Al-Khashman *et al*²¹ and Fagbote *et al*²² respectively. Total petroleum hydrocarbon content, nitrogen and phosphorus were done by methods describe by Khan *et al*²³, Okoye and Agbo²⁴, and Allen *et al*²⁵ respectively in their previous works.

Microbial analysis of abattoir and control soils: Microbial analysis of abattoir and Control soils were determined based on the method described by Rabah *et al*²⁶. Bacterial and Fungal Counts of the abattoir and Control soil samples were enumerated using pour plate method as described in details by Cappuccino *et al*²⁷. The results obtained were expressed as colony forming unit per gram of samples (CFU/g). The bacterial isolates which were maintained on agar slants were then characterized based on their cultural and morphological attributes and also on their responses to standard biochemical test as described by Cheesbrough M.²⁸ and identified as Brenner *et al*²⁹. Fungal isolates were characterized base cultural attributes, and identified by consulting various taxonomic books and monographs available on various groups of fungi³⁰.

Evaluation of trace metals pollution status in abattoir and Control soil samples: Trace metal pollution index: Trace metal pollution index denoted as MPI in the studied abattoir soils were calculated using equation (1) given below. Classifications of obtained MPI values were done according to Lacatusu³¹.

$$MPI = \frac{\text{Concentration of metal in studied soil}}{\text{Reference soil (control)}} \quad (1)$$

Degree of contamination (Cdeg) and Enrichment factor (EF): Degree of contamination (Cdeg) of each studied abattoir and Control soils was calculated using equation (2), while the enrichment factor (EF) of each metal was determined using equation (3) given below

$$Cdeg = \sum MPI \quad (2)$$

Where MPI = Metal pollution indexes for all the studied elements

$$EF = \frac{\left(\frac{M}{Fe}\right)^D}{\left(\frac{M}{Fe}\right)^C} \quad (3)$$

Where: D = Abattoir soils studied; C = Control; M = metal studied; Fe in the equation is used for normalization. The obtained Cdeg and EF values were then compared with different Cdeg and EF classifications given by Hakanson³², Rubio *et al*³³ and Zsefer *et al*³⁴.

Geo-accumulation index (Igeo): Igeo of Fe, Pb, Cd, Zn, Cu, Cr, and Ni in studied abattoir soils were determined using equation (4) below

$$Igeo = \log_2(Cn/1.5Bn) \quad (4)$$

Where: Cn = measured concentration of trace metal in studied samples, Bn = Concentration Control soil, Log 2 and 1.5 are constants described in Lu *et al*³⁵. The Igeo values obtained were then compared with the different Igeo models described in details by Huu *et al*³⁶.

Pollution load index (PLI): PLI of Fe, Pb, Cd, Zn, Cu, Cr, and Ni in abattoir soils were calculated using equation (5) given below.

$$PLI = (MPI_{Fe} \times MPI_{Pb} \times MPI_{Cd} \times MPI_{Zn} \times MPI_{Cu} \times MPI_{Cr} \times MPI_{Ni})^{\frac{1}{7}} \quad (5)$$

Where: MPI = Metal pollution indexes obtained using equation (1) above. The obtained PLI values were compared with the different PLI categories³⁷.

Results and discussion

Results for bulk density (gcm^{-3}), pH, and textural characteristics (quantity of sand, silt and clay) of abattoir and Control soils are presented in Table-1.

The results indicated varied textural characteristics, bulk density and soil pH among the studied abattoir soils and Controls. For the two abattoir soils, the percentage of sand, silt and clay were within the range of sand 50-58%, silt 8-15%, and clay 20-25%; while their Controls were 48-50% (sand), 15-17% (silt) and 18-24% (clay). The percentages of sand, silt and clay obtained for abattoir soils in this work agrees with 52-59% (sand), 10-16% (silt) and 25-30% (clay) reported by Edori *et al.* for abattoir soils from Port Harcourt, River State, Southern Nigeria but disagrees with 76-83% (sand), 1.5-2.0% (silt) and 13-23% (clay) reported by Ediene *et al.*³⁸ for abattoir soils from Calabar, Cross River State. All the abattoir soils studied and their Controls fell in the sandy-clay-loam (SCL) class of soil, with higher percentage of sand, followed by clay and then silt. Soil texture parameter was measure so as to reveal the physical properties of the soil such as water retention capacities, permeability, easy or toughness of tillage of the soil studied among other things. From these findings, the abattoir soils under investigation were seen to have potential of holding more water within the particle because of the high percentage of clay³⁹.

Results for bulk density (gcm^{-3}) and pH of abattoir soils and their Controls are presented in Table-1. The results indicate that

both parameters varied differently among the abattoir soils and the Control with bulk densities having ranges of 1.28-1.42 for the two abattoir soils and 1.14-1.17 for the Control. The result also reveals that the bulk densities of the abattoir soils were higher than those of the Control. The variations maybe as a result of differences in soil texture and organic matter content of abattoir soils and the Control. Also, Abattoir soil recorded greater percentages of sand than the Control soil and soils with higher percentage of sand is usually more prone to high bulk density¹⁷. The range of bulk density obtained in this work is in agreement with 1.16-1.81 gcm^{-3} reported by Olayinka *et al.*⁴⁰ for abattoir soils from Abeokuta, South western Nigeria, but in contrast with 1.50-1.65 gcm^{-3} reported by Chibuzor *et al.*⁴¹ for abattoir soils from Makurdi, Benue State, Nigeria. However, the values for bulk densities obtained in this work are considered suitable for crop production and also within the critical range^{42,43}.

The pH values for the two abattoir soils and their Controls recorded in this work were 4.78-5.30 and 4.60-4.74 respectively as indicated in Table-1. This parameter was determined because pH (acidity or alkalinity) plays a great role in determining nutrients availability and species of microorganism in soil⁴⁴. The results indicated that all the soils (abattoir and Control) studied were acidic in nature, with their Control soils showing higher acidities than the abattoir soils. However, the obtained ranges reported in this work are lower than 6.22-7.44 reported by Chukwu *et al.*⁴⁵ but are consistent with 4.99-6.73 obtained by Ubwa *et al.*⁵ in abattoir soils though with slight differences. Also, the observation of higher pH obtained for abattoir soils than in the Control soil obtained in this work is in line with the findings of Agbaire *et al.*⁴⁶. This could be attributed to biodegradable waste materials in studied abattoir soils which may lead to reduced anaerobic activities in these soils⁴⁷. Consequently, pH of soils impacted by abattoir wastes could be affected considerably. Although there is no acceptable standard for pH for an ideal soil for planting as it depends upon the type of crops, researches have shown that most minerals and nutrients are best available to plants in soil with a pH of between 6.5-6.8^{48,49}.

Table-1: Quantity (g/kg) of sand, silt and clay, textural class, bulk density and pH of abattoir soils and Control from Akwa Ibom State, Southern Nigeria.

| Parameters | Sampling points | | | | | |
|------------------------------------|-----------------|------|------|------|------|------|
| | IK1 | IK2 | IKC | EK1 | EK2 | EKC |
| Sand (%) | 58 | 55 | 48 | 56 | 52 | 50 |
| Silt (%) | 12 | 8 | 15 | 10 | 14 | 17 |
| Clay (%) | 23 | 22 | 18 | 25 | 20 | 24 |
| Texture | SCL | SCL | SCL | SCL | SCL | SCL |
| Bulk density (gcm^{-3}) | 1.42 | 1.28 | 1.14 | 1.38 | 1.36 | 1.17 |
| Soil pH* | 5.30 | 5.18 | 4.60 | 4.95 | 4.78 | 4.74 |

pH* (1:2.5 soil: water ratio); SCL – Sandy-clay-loamy soil; IK1-Ikot Ekpene abattoir soil 1; IK2-Ikot Ekpene abattoir soil 2; IKC-Ikot Ekpene control soil; EK1-Eket abattoir soil 1; EK2 - Eket abattoir soil 2; EKC - Eket control soil.

Table-2: Physicochemical properties of studied abattoir and Control soils.

| Parameters | Sampling points | | | | | |
|----------------------------------|-----------------|-------|-------|-------|-------|-------|
| | IK1 | IK2 | IKC | EK1 | EK2 | EKC |
| Temperature (°C) | 30.45 | 32.10 | 27.33 | 31.67 | 30.98 | 28.49 |
| EC (μS/cm) | 40.60 | 38.62 | 19.56 | 42.18 | 44.05 | 20.86 |
| Salinity (mgkg ⁻¹) | 18.00 | 22.00 | 10.50 | 15.00 | 26.06 | 13.15 |
| Moisture content (%) | 10.35 | 12.86 | 3.78 | 11.70 | 10.06 | 4.05 |
| TOM (%) | 8.65 | 7.94 | 4.37 | 7.27 | 8.42 | 4.64 |
| TOC (%) | 13.76 | 12.92 | 5.67 | 14.31 | 15.05 | 6.82 |
| CEC (Cmol/kg) | 28.63 | 26.80 | 20.84 | 25.11 | 26.95 | 21.67 |
| TPH (mgkg ⁻¹) | 8.06 | 7.15 | 2.61 | 11.72 | 13.63 | 4.52 |
| Nitrogen (%) | 0.04 | 0.06 | 0.01 | 0.09 | 0.10 | 0.02 |
| Phosphorus (mgkg ⁻¹) | 2.21 | 3.18 | 0.38 | 2.19 | 1.84 | 1.09 |

*EC–Electrical conductivity; TOM–Total organic matter, TOC–Total organic carbon, CEC–Cation exchange capacity, TPH–Total petroleum hydrocarbons; IK1-Ikot Ekpene abattoir soil 1; IK2-Ikot Ekpene abattoir soil 2; IKC-Ikot Ekpene control soil; EK1-Eket abattoir soil 1; EK2 - Eket abattoir soil 2; EKC - Eket control soil.

Results for the physicochemical properties of selected abattoir soils and their Controls are presented in Table-2. Temperature (°C) ranges for the two abattoir soils studied and their Controls were 30.45–32.10 and 27.33–28.49 respectively. The results also indicate that abattoir soils recorded varied temperatures and were higher than those of their respective Controls. The reasons for the differences in the temperatures among the abattoir soils may be attributed to factors such as variation in water content of the abattoir soils, soil relief and cover¹⁹. Temperature range for the abattoir soils obtained in this work is lower than (33.60–35.30)°C reported by Edori *et al*¹⁹ for abattoir soils from Port Harcourt, Rivers State, but higher than (18.80–21.43)°C reported by Osakia⁵⁰ for abattoir soils from Delta State, Nigeria. Temperature range for the Control soils recorded in this work is in agreement with 27.33–29.00°C reported by Rabah *et al*²⁶ for soils from different part of Southern Nigeria.

Levels of electrical conductivities (EC) in μS/Cm varied from 38.62 to 40.60 for abattoir soils obtained from IK, 42.18 to 44.05 for abattoir soils from EK and 19.56 -20.86 for the Control soils (Table-2). Ranges obtained for EC in studied abattoir soils are higher than 2.03–2.54 μS/Cm reported by Akan *et al*⁵¹ but lower than 60.00–110.00 μS/Cm obtained by Chukwu *et al*⁴⁵ in abattoir soils. Electrical Conductivity levels in studied abattoir soils were higher than values obtained at the Control soils, which is indicative of negative impact of abattoir wastes on studied soils. The findings of higher EC in abattoir soils than in Control are consistence with previous report⁵². This could be attributed to the low cation exchange capacity (CEC) of Control soil and variations in rates at which metallic salts and organic matter complexes are formed^{53,54}. Hence, EC of abattoir waste-impacted soils could be significantly affected by the

wastes. However, EC values recorded in abattoir soils in this study is within the permissible limit (below 100μS/Cm) stipulated by World Health Organization⁵⁵.

Ranges for salt content (mgkg⁻¹) of abattoir and Control soils as presented in Table-2 were 15.00-26.06 and 10.50-13.15 respectively. This result indicates that soil content of abattoir soils were higher than those of the Control soils. Although, salt content in soil are caused by natural factors such as weathering, continuous irrigation or pouring of wastewater (after washing of animal parts) on soil can also increase salt content of soil. This is so because almost all water contains some dissolve salts. Range for salinity of abattoir soils obtained in this study is lower than reported 29.00-59.00 mgkg⁻¹ by Edori *et al*¹⁹ and 475.05-667.88 mgkg⁻¹ by Chukwu *et al*⁴⁵ for abattoir soils from Port Harcourt, River State, Southern Nigeria. However, the abattoir soils recorded salinity values that are within permissible limit (200mgkg⁻¹) established for soil. The Low values of salinity recorded in abattoir soil is advantageous, since high salinity in soil usually leads to poor plant growth and lower soil microbial activity caused by osmotic stress and toxic ions¹⁵.

Moisture content in % varied from 10.35 to 12.86 for abattoir soils obtained from IK, 10.06 to 11.70 for abattoir soils from EK and 3.78 to 4.05 for the Control soils (Table-2). The variations in these values especially between the abattoir soils and the Control may be as a result of the effect of the abattoir effluent on soil. Ranges obtained for moisture content in studied abattoir soils are higher than 7.03-9.54% reported by Adesomoye *et al*⁵⁶ but lower than 19.01-21.07% obtained by Edori *et al*¹⁹ and 17.91-19.50% obtained by Olayinka *et al*⁴⁰ in abattoir soils. The findings of higher moisture content in abattoir soils than in

Control are consistence with the report of Onweremadu *et al*⁵². This observation can be explained by the fact that in ruminants, the first stomach or paunch contains undigested materials or paunch manure. The paunch manure which is disposed on the soil could have a moisture content of about 88%¹⁰.

Results in Table-2 indicate ranges for total organic matter (TOM) for IK, EK abattoir soil and Control soil as 7.94-8.65%, 7.27-8.42% and 4.37-4.64 respectively. These ranges are higher than 0.69-7.42% reported by Yahaya *et al*³ for abattoir soils, and is inconsistent with values obtained by Neboh *et al*⁵⁷ but lower than 5.57-24.13% obtained by Ubwa *et al*⁵. Values of OM obtained in studied abattoir soils were also higher than values at Control which is consistent with report by Owagboriaye *et al*⁵⁸ and Oyedele *et al*⁵⁹. This disparity may be attributed to the absence of biodegradable wastes at Control site thereby indicating that; waste materials from abattoir may increase the OM of soil significantly. Also, the faeces of livestock have been observed to consist of undigested food which hitherto will increase the OM content of abattoir soil¹⁰. However, soil organic matter usually act as a “storehouse or reservoir” for most metals hence it can influence their availability in soil either positively or negatively⁴⁶.

Total organic carbon (TOC) results indicated ranges of 12.92-15.05% for the abattoir soils and 5.67-6.82% for the Control soils. The Control soils recorded lower TOC than the abattoir soils. This could be as a result of higher organic matter levels of abattoir soil. This observation corroborates with the reports of Yahaya *et al*³ and Alorge⁶⁰ in their respective studies. Results obtained for studied abattoir soils in this study are lower than 6.1-7.6% reported by Ediene *et al*³⁸ for abattoir soil from Calabar Metropolis, Cross River State Nigeria, but higher than 12.68-30.02% reported by Edori *et al*¹⁹. The differences in the reported values of total organic carbon and those earlier reported for abattoir soils may be due to the rate of decomposition and composting of animal wastes such as dung, body part, blood, bones etc³. Organic carbon content in soil plays a vital role in soil development, fertility, and moisture availability in soil.

Cation exchange capacity (CEC) of studied abattoir soils varied from 26.80 to 28.63 Cmolkg⁻¹ for IK abattoir soils, 25.11 to 26.95 Cmolkg⁻¹ for EK abattoir soils, and 20.84 to 21.67 Cmolkg⁻¹ for the Control soils (Table-2). The result also shows that the respective abattoir soils recorded higher CEC than the Control soils. This is may be due to higher total organic matter content of the abattoir soils than in the Control soils including the clay content of the soil⁶¹. Higher content of CEC in abattoir soils than in the Control soils obtained in this study agrees with the findings of Iwegbue *et al*⁶². Also, the obtained ranges of CEC in abattoir soils are higher than 12.54-16.84 Cmolkg⁻¹ reported by Neboh *et al*⁵⁷ in abattoir soils. CEC measures soil's ability to hold positively charged ions and is very vital to plant as it influences soil structure stability, nutrient availability, soil hydrogen concentrations (pH), and the soil's reaction to

fertilizers and other ameliorants⁶³. Although most crops do well in soil with low CEC, but vegetables and other productive food crops like vegetables are perform best in soil with moderate to high CEC⁶⁴. This study thus reveals the effect abattoir wastes have on CEC of soils laden with it (abattoir wastes).

Results presented in Table-2 reveals that total petroleum hydrocarbons (TPH) content of the abattoir soils ranged from 7.15-8.06 mgkg⁻¹ for abattoir soil samples from IK, 11.72-13.63 mgkg⁻¹ for abattoir soils from EK and 2.61-4.52 mgkg⁻¹ for the Control soils. For the abattoir soils, ranges obtained in this study is within the reported range of 11.37-27.68 mgkg⁻¹ by Edori *et al*¹⁹ for abattoir soils from Port Harcourt, River State, Nigeria. EK abattoir soils recorded higher TPH content than IK abattoir soils as shown in the Table-3. This is because EK hosts some oil companies like Exxon Mobil PLC, while there is no such in IK. For the Control soils, the range obtained in this study is lower than reported 3400-6800 mgkg⁻¹ by Osuji *et al*⁶⁵, for abattoir soils from Owaze, Abia State, and 581.02 mgkg⁻¹ reported by Okoro *et al*⁶¹. The large variation in the reported values of TPH content for the Control soils in this study and those earlier reported is as a result of lesser frequency of crude oil spill in the Control site used for this study. Areas of frequent crude oil spillage are expected to have higher TPH values than those of sparsely crude oil spillage. However, the lower concentrations of TPH in both abattoir and Control soils in this study is still a source of concern as TPH have been reported as a contaminant in any environment due to its toxicity to humans and other environmental receptors⁶⁶.

Total nitrogen content (%) (both as NH₄⁺ - N and NO₃⁻ - N) and phosphorus (mgkg⁻¹) of the abattoir soils and Control soils are presented in Table-2. For Nitrogen, ranges are 0.04 – 0.06; 0.09 – 0.10 and 0.01 – 0.02 for IK, EK and Control soils respectively, while 2.21–3.18, 1.84–2.19 and 0.38-1.09 for IK, EK and the Control soils respectively. The variations between nitrogen and phosphorus contents among the abattoir and Control soils obtained in this study may be as a result of varied amounts of nitrogenous compounds in the abattoir impacted soils and the Control. The results also show that the abattoir soils recorded higher content of Nitrogen than the Control soils. This finding agrees the report of Ediene *et al*³⁸ who also reported higher nitrogen content for abattoir soils from Calabar, Cross River State than in the Control. Nitrogen content in abattoir soil obtained in this study is lower than 0.48 – 3.01% reported by Edori *et al*¹⁹ and 0.18 - 0.65% reported by Ediene *et al*³⁸ for abattoir soils, but higher than (0.008-0.009%) reported by Abubakar⁶⁷ for abattoir soils from Yola metropolis, Adamawa State, North Eastern Nigeria. For the Control soil, the obtained range is lower than 0.08–0.09% reported by Ediene *et al*³⁸. For phosphorus, both abattoir soils recorded higher phosphorus content than the Control soils. This result also agrees with the report of³⁸. This could be explained by the higher pH (less acidity) values and higher organic matter of the abattoir soils as indicated in Table-2. Range of phosphorus obtained in this study is consistent with reported 2.46 – 3.61

mgkg⁻¹ by Abubakar⁶⁷, though lower than 0.005-0.007 mgkg⁻¹ reported by Rabah *et al*²⁶ for abattoir soils obtained from Sokoto State, Nigeria. Both nitrogen and phosphorus contents of abattoir soils obtained in this study all fell within the permissible limits (nitrogen < 40% and phosphorus < 40 mgkg⁻¹) stipulated by World Health Organization⁵⁵ for soils. Although, nitrogen and phosphorus are needed in soil by plants since the first is a building block of protein, nucleic acid and other cellular constituents which are essential to all forms of life, the later is a component of the complex nucleic acid structure of plants, which regulates protein synthesis in plants⁶⁸. However, excess nitrogen and phosphorus in soil usually causes plants to mature too rapidly in addition to reducing Zn, Cu and Fe availability in soil. Consequently, the moderate levels of nitrogen and phosphorus in the abattoir soils under investigation is plausible in lieu of its usage for planting of crops⁴⁵.

Results for essential elements and trace metal levels of abattoir soils and Control soils are presented in Table-3. The results indicated significant variations in the levels of all essential and trace metals investigated for the abattoir and Control soils. Results for potassium (K) indicated the following ranges: 0.81-0.93 Cmolkg⁻¹ for abattoir soils from IK, 0.68 – 0.74 Cmolkg⁻¹ for EK and 0.17-0.26 Cmolkg⁻¹ for the Control. The reason for this variation may be as a result of the high moisture content of the abattoir soils as indicated in Table-3. Levels of potassium in the Control soils were lower than those of the abattoir soils and this finding corroborates with the report of Neboh *et al*⁵⁷ who also reported similar findings. Ranges of potassium obtained in this study for the abattoir soils are consistent with the reports of Sumayya *et al*⁶⁹ and Neboh *et al*⁵⁷ in their respective studies. Potassium has many different roles in the soil relative to plants. It is involve in photosynthesis as it regulate the opening and closing of stomata, regulate carbon (IV) oxide, triggers activation of enzymes and its essential for production of

adenosine triphosphate (ATP)⁶⁴. For other exchangeable bases (Sodium, calcium and magnesium), abattoir waste had significant influences on their levels in the studied abattoir soils.

From the results presented in Table-3, the abattoir soils recorded highest value of Na, Ca and Mg than in Control soils. In the case of Na, ranges were 0.42-0.51molkg⁻¹ for samples from IK, and 0.53 -0.67molkg⁻¹ for samples from EK. Ca and Mg recorded ranges of 1.83-2.04 molkg⁻¹ and 2.08-3.92molkg⁻¹ for IK, while EK were 1.65-1.83 and 2.19-2.67 molkg⁻¹. Ranges of Na obtained in this study for abattoir soils is higher than reported 0.1-0.12 molkg⁻¹ by Ediene *et al*³⁸, but lower than 2.24-2.47 molkg⁻¹ reported by Abubakar⁶⁷ for abattoir soils obtained from Yola Metropolis, Adamawa State, Nigeria. Ca and Mg ranges obtained for abattoir soils in this study are higher than 0.48 – 0.50 mol/kg Ca and 0.51 – 0.77 molkg⁻¹ Mg reported by Abubakar⁶⁷ but lower than 12.6–15.6 molkg⁻¹ Ca and 4.06 – 9.80 molkg⁻¹ reported by Ediene *et al*³⁸. The Control soil samples recorded lower levels of Na, Ca and Mg than the studied abattoir soils. These results agree with previous reports of Neboh *et al*⁵⁷ and Ediene *et al*³⁸ although at variance with the report of Abubakar⁶⁷, who reported lower Ca level in abattoir soils than in the Control soils. Several factors affect the levels of exchangeable bases in soil and this includes texture and soil organic matter, CEC and soil moisture content⁶⁴. Exchangeable bases such as Na, Ca and Mg are important in soil because they are involve in translocation of carbohydrates and nutrient within plants, cell growth, are component of chlorophyll for photosynthesis, protein synthesis and energy transfer within plants. Although, there are no limits to the amount of these bases in soil, higher levels of Na usually causes dispersion of fine particle of soils into pores thereby reducing water penetration and blocking plant root access. Higher levels of Ca in soil reduces uptake of other cation nutrients⁵⁷.

Table-3: Essential elements and trace metals levels of selected abattoir and Control soils.

| | Unit | Sampling Points | | | | | |
|--------------------|-------------|-----------------|--------|--------|--------|--------|--------|
| | | IK1 | IK2 | IKC | EK1 | EK2 | EKC |
| Essential elements | K (mol/kg) | 0.93 | 0.81 | 0.26 | 0.74 | 0.68 | 0.17 |
| | Na (mol/kg) | 0.42 | 0.51 | 0.16 | 0.67 | 0.53 | 0.32 |
| | Ca (mol/kg) | 1.83 | 2.04 | 1.23 | 1.65 | 1.83 | 0.98 |
| | Mg (mol/kg) | 2.08 | 3.92 | 0.83 | 2.67 | 2.19 | 1.03 |
| Trace metals | Fe (mg/kg) | 643.45 | 604.76 | 548.10 | 611.04 | 665.10 | 562.82 |
| | Zn (mg/kg) | 19.23 | 21.05 | 15.09 | 24.13 | 18.56 | 12.15 |
| | Cd (mg/kg) | 0.35 | 0.47 | 0.17 | 0.46 | 0.52 | 0.23 |
| | Cu (mg/kg) | 16.82 | 14.94 | 4.94 | 20.92 | 16.30 | 3.56 |
| | Pb (mg/kg) | 0.73 | 0.66 | 0.32 | 1.01 | 0.89 | 0.37 |
| | Cr (mg/kg) | 0.32 | 0.26 | 0.05 | 0.18 | 0.22 | 0.08 |
| | Ni(mg/kg) | 9.73 | 11.47 | 6.19 | 10.21 | 8.84 | 4.23 |

*IK1-Ikot Ekpene abattoir soil 1; IK2-Ikot Ekpene abattoir soil 2; IKC-Ikot Ekpene control soil; EK1-Eket abattoir soil 1; EK2 - Eket abattoir soil 2; EKC - Eket control soil.

Results in Table-3 indicate that, Fe varied between (604.76–643.45mgkg⁻¹) for IK abattoir soils, (611.04– 665.10mgkg⁻¹) for EK abattoir soils, and (548.10–562.82mgkg⁻¹) for the Control soils. Levels of Fe obtained for both abattoir soils in this study is lower than 2569.00–4130.00mgkg⁻¹ reported by Yahaya *et al*³, but higher 59.36– 81.70mgkg⁻¹ obtained by Simeon *et al*⁷⁰. Also, from the results, abattoir soils recorded higher levels of Fe than the Control soils and this is in agreement with the report of Neboh *et al*⁵⁷. This is indicative of additional source of Fe in studied abattoir wastes-impacted soils. EK2 sample recorded highest Fe level, while IK2 recorded lowest for all the samples studied. Results obtained in this study revealed direct relationship between activities at abattoir and Fe accumulation in studied abattoir soils. However; levels of Fe in both studied abattoir soils and Control are higher than 400.00mgkg⁻¹ recommended by Federal Environmental Protection⁷¹ for Nigerian soils. This confirms that Nigerian soils have elevated levels of Fe as clearly shown by their reddish nature. Nevertheless, the availability of Fe in soil for plant uptake may not be guaranteed as Fe oxides (the major form of Fe in soil) are highly insoluble in soil⁷².

Results obtained showed that concentrations of Zn in studied abattoir soils varied between (19.23–21.05mgkg⁻¹) for IK abattoir soils, (18.56–24.13mgkg⁻¹) for EK abattoir soils, and (12.15–15.09mgkg⁻¹) for the Control soils. Ranges of Zn obtained in this study are lower than 1.302–5.2362mgkg⁻¹ reported by Ubwa *et al*⁵ in abattoir soils, but lower than 50.91–92.50mgkg⁻¹ obtained by Yahaya *et al*³ and 171.93mgkg⁻¹ obtained by Olayinka *et al*⁴⁰. Levels of Zn reported in studied abattoir soils were relatively higher than values obtained at the Control site. This is in agreement with the report of Neboh *et al*⁵⁷ who also reported a higher level of Zn in abattoir soils than in Control soils. However; the obtained ranges are lower than 140.0 mgkg⁻¹ limit by Federal Environmental Protection⁷³ for Nigerian soils. Nevertheless, since toxicity of metal may not be identified by total metal concentration alone its availability may not be established. Also, lower levels of Zn obtained in this study for Zn when compared to the permissible limit in soil set by FEPA is plausible because Zn though very essential in soil, it is needed by plant in trace amount.

For Cd, results indicated the following ranges: (0.35 – 0.47mgkg⁻¹) for IK abattoir soils, (0.46 -0.52mgkg⁻¹) for EK abattoir soils, and (0.17–0.23mgkg⁻¹) for Control soils. For the abattoir soils, Cd showed highest abundance in abattoir soil from EK2 and least in abattoir soil from IK1, while Control soils recorded lower levels of Cd than the abattoir soils studied. Ranges for Cd obtained in this study agrees with 0.43–0.71 mgkg⁻¹ obtained by Chukwu *et al*⁴⁵ for abattoir soils from Obiaakpor Area, River State, Nigeria, but higher than 0.25 mgkg⁻¹ reported by Olayinka *et al*⁴⁰. However, levels of Cd obtained were below 2.0 -3.0 mgkg⁻¹ permissible limit stipulated for soil by Federal Environmental Protection⁷³. This shows that studied abattoir soils were not overloaded with Cd even though other parameters may need to be evaluated to really ascertain the status of the abattoir soils as it relates to Cd.

Results in Table-3 indicate ranges for Cu as 14.94–16.82mgkg⁻¹, 16.30–20.92mgkg⁻¹ and 3.56–4.94mgkg⁻¹ for IK, EK abattoir soils and Control soils respectively. Ranges of Cu obtained for abattoir soils are higher than reported 0.05–1.7mgkg⁻¹ by Osu *et al*⁶⁵ for abattoir soils in Umuahia, Nigeria but lower than reported 36.46–40.60mgkg⁻¹ obtained by Owagboriaye *et al*⁵⁸ in Abeokuta, Nigeria. The highest concentration of Cu was recorded in samples from EK1 and lowest in those from IK1. Concentrations of Cu in studied abattoir soils were higher than values obtained at the Control Soils. This indicated availability of Cu-containing waste materials in studied abattoir waste-impacted soils. This is consistent with the report of Chukwu *et al*⁴⁵ for abattoir soils. The obtained Cu values are also lower than 36.0mgkg⁻¹ stipulated by FEPA for Nigerian soils. Nonetheless, bioavailability and toxicity of Cu could not be confirmed based on total concentration alone.

Results obtained for total Pb in studied abattoir soils indicated a range of 0.66–0.73mgkg⁻¹ and 0.89–1.01mgkg⁻¹ for IK and EK abattoir soils (Tables-3). Levels of Pb obtained in studied abattoir soils are lower than reported 7.17–12.50mgkg⁻¹ by Chukwu *et al*⁴⁵ for abattoir soils from Obio Akpor, River State Nigeria but higher than reported 0.18 – 0.83mgkg⁻¹ by Ubwa *et al*⁵ except for IK abattoir soils. Highest level of Pb was reported in samples from EK1 while lowest Pb concentration was obtained in IK2 abattoir soil. Concentration of Pb in studied abattoir soils were higher than values obtained in the Control site revealing negative impact of abattoir wastes on Pb levels in studied soils. However; levels of Pb obtained were below 85.00mgkg⁻¹ recommended by Federal Environmental Protection⁷³ for soil in Nigeria.

Levels of Cr and Ni in studied abattoir soils (IK and EK) ranged from 0.26–0.32mgkg⁻¹ and 0.18–0.22mgkg⁻¹ for Cr and 9.73–11.47mgkg⁻¹ and 8.84–10.21mgkg⁻¹ for Ni (Tables-3). The ranges for Cr are lower than reported 4.25–5.86mgkg⁻¹ by Chukwu *et al*⁴⁵ but higher reported 0.0717–0.1358mgkg⁻¹ by Ubwa *et al*⁵ in abattoir soils. For Ni, Levels of Ni obtained are higher than 2.160–4.690mgkg⁻¹ reported by Simeon *et al*⁷⁰ but lower than 33.50–107.13mgkg⁻¹ recorded for Ni in abattoir soils by Yahaya *et al*³. The highest Cr level was obtained in abattoir soil from IK1, while the lowest concentration was in EK1 abattoir soil. For Ni, highest Ni level was obtained in abattoir soil from IK2, while the lowest level was obtained in EK2 abattoir soil. Levels of Cr and Ni in abattoir soils were higher than levels obtained in soil from the Control site and are in agreement with findings reported by Chukwu *et al*⁴⁵ in abattoir soils.

However, values of Cr and Ni obtained in studied abattoir soils are much lower than 100.0mgkg⁻¹ Cr and 35.0mgkg⁻¹ Ni limits in soil by Federal Environmental Protection⁷³. Nevertheless, pollution status of these metals may not be ascertained using information from total concentration, but lower levels of Cr and Ni obtained in this study are significant especially since abattoir soils studied are already used by farmers in planting crops.

Trace metals pollution status in abattoir and Control soil samples: To ascertain the pollution status of the studied abattoir soils, MPI, Cdeg, trace metals enrichment factor (EF), geo-accumulation index (Igeo) and metal pollution load (MPL) were evaluated. Metal pollution index (MPI) was used to differentiate between contamination and pollution levels in studied abattoir soils as indicated in equation (1) above⁷⁴. The different categories of MPI for each of the trace metals as indicated in Table 5 shows that Fe recorded MPI values ranging from 1.09–1.53 for all abattoir soils under investigation indicating slight pollution of the studied abattoir soils by Fe. Consequently, it can be predicted that the level of Fe in the studied abattoir soils can impose a negative impacts on the surrounding environment of the study area. However, since Fe in soil is usually characterized with low bioavailability factor hence; the consequences may not be alarming. Results in Table-4 also indicates that MPI values for Zn was in slight pollution (1- 2) category, while Cd, Cu, Pb, Cr, and Ni were in moderate pollution (2.1–4.0) category except for abattoir soils from EK1 and EK2 that were in severe pollution (4.1-8.0) category for Cu and IK1 and IK2 for Cr. This means that Zn, Cd, Cu, Pb, Cr and Ni have slightly polluted the studied abattoir soils and is expected to affect the soil, plants and the studied environment negatively.

Enrichment factors (EF) of metals were calculated for the abattoir soils using the continental crust average model (Table-4). Fe exhibited an EF value of 1.00 in all the abattoir soils studied indicating that a greater proportion of Fe may have emanated from natural soil forming processes³⁴. EF for Zn, Cd, Cu, Pb, Cr and Ni for all the abattoir soils studied were greater

than 1.0 indicating that these trace metals were from anthropogenic source³⁴. Results for Igeo of trace metals in studied abattoir soils are presented in Table-4. Results obtained showed the following ranges: 0.26–0.39 Zn, 0.40–0.55 Cd, 0.61–1.18 Cu, 0.41–0.56 Pb, 0.45–1.28 Cr and 0.31–0.48 Ni for all the abattoir soils studied. From these results, the metals were in the 0–1 class (unpolluted – moderately polluted) according to Huu et al³⁶. Cu In abattoir soil from EK1 and Cr in abattoir soil from IK1 were in the 1–2 class (moderately polluted).

Results for the degree of contamination (Cdeg) of the four (4) studied abattoir soils investigated are presented in Figure-1. From the results, Cdeg values were 18.10, 17.38, 18.33 and 17.15 for IK1, IK2, EK1, and EK2 respectively. The varied Cdeg reported in this study by the abattoir soils may be attributed to the volume of abattoir wastes and abattoir activities in each of these sites. From the Cdeg results, it can therefore be deduced that the abattoir soils were considerably contaminated ($16 < Cdeg < 32$) based on the model predicted by Hakanson³².

Results obtained for PLI for the four abattoir soils examined are indicated in Figure-1. PLI values were found to range from 2.17 for abattoir soil from IK1 to 2.33 for abattoir soil from EK1. Thus, PLI values of all the studied abattoir soils were within the heavy pollution ($2 < PLI < 3$) category according to Tomlinson et al³⁷. These PLI values obtained in this study further confirm findings by degree of contamination of studied abattoir soils. This work has therefore revealed the negative impact of abattoir wastes on underlying soils with regards to metal accumulation.

Table-4: MPI, EF and Igeo of trace metals in studied abattoir soils.

| | | Fe | Zn | Cd | Cu | Pb | Cr | Ni |
|------|-----|------|------|------|------|------|------|------|
| MPI | IK1 | 1.17 | 1.20 | 2.08 | 3.40 | 2.28 | 6.40 | 1.57 |
| | IK2 | 1.10 | 1.39 | 2.76 | 3.02 | 2.06 | 5.20 | 1.85 |
| | EK1 | 1.09 | 1.98 | 2.00 | 5.87 | 2.73 | 2.25 | 2.41 |
| | EK2 | 1.53 | 1.53 | 2.26 | 4.58 | 2.41 | 2.75 | 2.09 |
| EF | IK1 | - | 1.09 | 1.75 | 2.90 | 1.94 | 5.45 | 1.33 |
| | IK2 | - | 1.26 | 2.51 | 2.74 | 1.87 | 4.71 | 1.68 |
| | EK1 | - | 1.83 | 1.84 | 5.41 | 2.51 | 2.07 | 2.33 |
| | EK2 | - | 1.29 | 1.01 | 3.87 | 2.04 | 2.33 | 1.77 |
| Igeo | IK1 | - | 0.26 | 0.41 | 0.68 | 0.46 | 1.28 | 0.31 |
| | IK2 | - | 0.28 | 0.55 | 0.61 | 0.41 | 1.04 | 0.37 |
| | EK1 | - | 0.39 | 0.40 | 1.18 | 0.56 | 0.45 | 0.48 |
| | EK2 | - | 0.31 | 0.45 | 0.92 | 0.48 | 0.55 | 0.41 |

*MPI – Metal pollution index; EF – Enrichment factor; Igeo – Geo-accumulation index; IK1-Ikot Ekpene abattoir soil 1; IK2-Ikot Ekpene abattoir soil 2; IKC-Ikot Ekpene control soil; EK1-Eket abattoir soil 1; EK2 - Eket abattoir soil 2; EKC - Eket control soil.




Figure-1: Degree of contamination (Cdeg) and pollution load index (MPL) of trace metals in studied abattoir soils.




Figure-2: Microbial Load (CFU/g) of abattoir soils obtained from Ikot Ekpene and Eket region of Akwa Ibom State.

Microbial loads of abattoir soils: Figure-2 shows the microbial load (\log_{10} CFU/g) of soil samples. The result shows the density of the culturable bacterial community present in 1g of samples obtained from the four (4) abattoir locations ranges from 6.41 ± 0.43 - $7.91 \pm 0.58 \log_{10}$ CFU/g for total heterotrophic bacterial count and 4.94 ± 0.26 to $5.79 \pm 0.34 \log_{10}$ CFU/g for fungal count. Among the four (4) locations in which samples were obtained for the study, IK2 had the highest heterotrophic bacterial densities of $7.91 \pm 0.58 \log_{10}$ CFU/g while IK1 had the highest fungal count of $5.79 \pm 0.34 \log_{10}$ CFU/g.

Cultural and biochemical characteristics of microbial isolates: The cultural and biochemical properties of the microbial isolates are presented in Table-5 and 6. Six (6) species of bacteria were obtained from the soil samples using the aerobic culture techniques. These were *Klebsiella*, *Micrococcus*,

Pseudomonas, *Bacillus*, *Escherichia* and *Enterobacter* species. While the two (2) fungal species isolated were *Aspergillus* and *Penicillium* species.

Occurrence and Distribution of microbial isolates within the samples and location: The distribution of microbial isolates within the soil samples is shown in Table-7 and 8. *Micrococcus luteus*, *Pseudomonas aeruginosa* and *Bacillus subtilis* 5(100%) had the highest frequency of occurrence, while *Enterobacter cloacae* and *Klebsiella pneumoniae* had the least 2(40%). For the fungal isolates, *Aspergillus niger* 5(100%) had the highest frequency of occurrence, while *Penicillium frequentans* had the least 3(40%) frequency of occurrence. Also, samples taken from Ikot Ekpene region were more contaminated than those taken from Eket abattoir soils.

Table-5: Morphological and Biochemical characteristics Bacterial of Isolates.

| Isolates | G.R | Shape | Motility | Catalase | Starch hydrolysis | Oxidase | Indole | Citrate | MR | VP | Glucose | Lactose | Mannitol | Probable Organism |
|----------|-----|-------|----------|----------|-------------------|---------|--------|---------|----|----|---------|---------|----------|-------------------------------|
| 1 | - | R | - | + | - | - | + | + | - | + | AG | AG | - | <i>Klebsiellapneumoniae</i> |
| 2 | + | S | - | + | - | - | - | - | | | - | - | - | <i>Micrococcus luteus</i> |
| 3 | - | R | + | + | - | - | - | + | - | + | AG | AG | - | <i>Enterobactercloacae</i> |
| 4 | + | R | - | + | + | - | + | + | - | - | - | - | - | <i>Bacillus polymyxa</i> |
| 5 | + | R | - | + | + | - | - | - | - | - | - | - | - | <i>Bacillus subtilis</i> |
| 6 | - | R | + | + | - | + | - | + | - | - | A | - | A | <i>Pseudomonas aeruginosa</i> |
| 7 | - | R | + | + | - | - | + | - | + | - | AG | AG | - | <i>Echerichia coli</i> |

Key: G.R = Gram Staining; + = Positive; - = Negative; R = Rod; S = Spherical; A = Acid only; AG = Acid and Gas produced.

Table-6: Colonial and morphological characteristics of fungal (mould) isolates.

| Isolates | Colonial Morphology | Somatic cell type | Type of Hyphae | Asexual spores | Special Reproductive Structure | Conidia Head | Vesicle shape | Probable Fungi |
|----------|--------------------------------------|-------------------|------------------|-----------------|--------------------------------|----------------------------|---------------|--------------------------------|
| 1 | Compact white with dark basal colour | Filamentous | Septate mycelium | Globose conidia | Conidiospore | Globose | Subglobose | <i>Aspergillusniger</i> |
| 2 | Whitish, yellowish to grey mycelium | Filamentous | Septate mycelium | Globose conidia | Conidiospore | Sub-globose to ellipsoidal | - | <i>Penicillium frequentans</i> |

Table-7: Occurrence and distribution of bacterial isolates in abattoir and Control soil samples.

| Isolates | Sample Points | | | | EK2 | % Occurrence |
|-------------------------------|---------------|-----|-----|-----|-----|--------------|
| | Control | IK1 | IK2 | EK1 | | |
| <i>Klebsiella puemoniae</i> | - | + | + | - | - | 2(40%) |
| <i>Micrococcus luteus</i> | + | + | + | + | + | 5(100%) |
| <i>Enterobacter cloacae</i> | - | + | + | - | - | 2(40%) |
| <i>Pseudomonas aeruginosa</i> | + | + | + | + | + | 5(100%) |
| <i>Escherichia coli</i> | - | + | + | + | + | 4(80%) |
| <i>Bacillus polymyxa</i> | + | + | + | - | - | 3(60%) |
| <i>Bacillus subtilis</i> | + | + | + | + | + | 5(100%) |

*Key: + = Present; - = Absent.

Table-8: Occurrence and distribution of fungal isolates in abattoir and Control soil samples.

| Isolates | Sample Points | | | | | % Occurrence |
|------------------------------|---------------|-----|-----|-----|-----|--------------|
| | Control | IK1 | IK2 | EK1 | EK2 | |
| <i>Aspergillusniger</i> | + | + | + | + | + | 5(100%) |
| <i>Penicillumfrequentans</i> | + | + | + | - | - | 3(60%) |

*Key: + = Present; - = Absent.

The significant ($p=0.05$) increase in microbial loads encountered in the abattoir soil against the control is not surprising and it could be directly linked to the impacted abattoir wastes disposed on the studied land. This is because abattoir wastes contained substances that may perhaps serve as a utilizable source of nutrient and encourage rapid multiplication by microorganisms. These findings is consistent with the report of Adesemoye *et al*⁵⁶ and Rabah *et al*²⁶ who independently reported similar increase in microbial load of soil samples contaminated with abattoir effluents. Eket (EK1 and EK2) abattoir soils were found to harbor less number and species of microorganisms. This could be because the region is well known for oil exploration activities and is expected to harbor more of microorganisms that could survive in soil contaminated with hydrocarbon. The presence and abundance of species of *Bacillus* and *Micrococcus*, *Aspergillus* and *Penicillium* observed in both the abattoir and Control soils is not shocking as *Bacillus*, *Micrococcus*, *Aspergillus* and *Penicillium* microorganisms are local to soil surroundings and usually endure and thrive especially during carbon and nitrogen sources influx in the soil⁷⁵. However, presence of *Pseudomonas aeruginosa*, *E. coli*, *Enterobacter cloacae* and *Klebsiella pneumoniae* in the abattoir soil may be attributable to large amounts of feces in abattoir wastes since these microorganisms are well known flora of fresh beef. Their presence in the abattoir soil samples is indicative of recent faecal pollution as they are mostly indicator organisms. Similar findings were reported by Rabah *et al*²⁶. A good number of fungal isolated from abattoir soils in this work lives in soil and are known for their spoiling roles in beef^{75,76}. This in effect shows that the studied (abattoir) soils may have been polluted with abattoir wastes from abattoir activities in the studied sites.

Conclusion

The results of this study have shown the physicochemical characteristics, total essential and trace metal levels, total heterotrophic bacterial and fungal loads of abattoir and Control soils from Ikot Ekpene and Eket Local Government Areas of Akwa Ibom State, Nigeria. Metal pollution index (MPI), enrichment factor (EF), geo-accumulation index (Igeo), degree of contamination (Cdeg) and pollution load index (PLI) of trace metals have also been calculated using empirical pollution models. Essential elements and trace metal levels were higher in abattoir soils than in Control though were within permissible limit in soil except for Fe. Also, microbial results revealed a significant increase in the number and diverse of

microorganisms in the abattoir soils, some being disease causing, but are more often than not indicators of recent faecal pollution in the soil impacted with abattoir wastes. This study, therefore, concludes that soil impacted with abattoir wastes is richer in plant nutrients and can be exploited for growing of crops. But it is advised that routine checks be conducted to forestall trace metals accumulation above safe levels in these soils.

References

1. Nwuche C.O. and Ugoji E.O. (2010). Effect of Co-existing Plant Species on Soil Microbial Activity under Heavy Metal Stress. *International Journal of Environmental Science and Technology*, 7(4), 697-704.
2. Daniel L.A. and Braide S.A. (2004). The Effect of Oil Spillage on a Cultivated Wet-land of the Niger Delta. *Journal of Environmental Science*, 2, 153-158.
3. Yahaya M.I., Mohammad S. and Abdullahi B.K. (2009). Seasonal Variations of Heavy Metals Concentration in Abattoir Dumping Site Soil in Nigeria. *J. Appl. Sci. Environ. Manage.*, 13(4), 9-13.
4. Hornby A.S. (2006). Oxford Advanced Learner's Dictionary of current English. 7th edition, 1381.
5. Ubwa S.T., Ato G.H., Offem J.O., Abah J. and Asemave K. (2013). Effect of Activities at the Gboko Abattoir on Some Physical Properties and Heavy Metals Levels of Surrounding Soil. *International Journal of Chemistry*, 5(1), 49-57.
6. Red Meat Abattoir Association (2010). Waste Management-Red Meat Abattoir. Retrieved from SON (2003). Standard Organization of Nigeria Safe Drinking Water Regulation. <http://www.docstoc.com/docs/103302144/Waste-Management-%EE%9F% A6-Red-Meat-Abattoirs>.
7. Steffen L.C., Roberts H.Y. and Kirsten (1989). Water and waste-water management in the red meat industry. 36, WRC Report No. 145 TT41/89. WRC, Pretoria.
8. Kruslin E., Hodel C.M. and Schurgast H. (1999). Progress in Diagnosis of Chronic Toxic Metal Poisoning by Hair Analysis. *Toxicol. Lett.*, 88, 84-90.

9. Jukna C., Jukna V. and Siugzdaite J. (2006). Determination of heavy metals in viscera and muscles of cattle. *Bulgarian Journal of Veterinary Medicine*, 9(1), 35-41.
10. Ezeoha S.L. and Ugwuishiwu B.O. (2011). Status of Abattoir Wastes Research in Nigeria. *Nigerian Journal of Technology*, 30(2), 143-148.
11. Nwachukwu M.I., Akinde S.B., Udujih O.S. and Nwachukwu I.O. (2011). Effect of Abattoir Wastes on the Population of Proteolytic and Lipolytic Bacteria in a Recipient Water Body (Otamiri River). *Global Research Journal of Science*, 1, 40-42.
12. Magaji J.Y. and Chup C.D. (2012). The Effects of Abattoir Waste on Water Quality in Gwagwalada-Abuja, Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 5(4), 542-549.
13. Raymond C.L. (1977). Pollution Control for Agriculture. Academic Press Inc. New York, 45-58.
14. Saidu M. and Musa J.J. (2012). Impact of Abattoir Effluent on River Landzu, Bida, Nigeria. *Journal of Chemical, Biological and Physical Sciences*, 2(1), 132-136.
15. Chen T., Liu X., Li X., Zhao K., Zhang J., Xu J., Shi J. and Dahlgren R.A. (2009). Heavy Metal Sources Identification and Sampling Uncertainty Analysis in a Field-scale Vegetable Soil of Hang Zhou, China. *Environmental Pollution*, 157(3), 1003-1010.
16. Ibitoye A.A. (2006). Laboratory Manual on Basic Soil Analysis. Foladave Nigeria Limited, Ibadan, 16-36.
17. Olofunmi O.E. and Alli A.A. (2016). To Determined Variability of Soil Physical and Chemical Properties of a Typical Land Disposal Catfish Effluent in South Western Nigeria. *International Journal of Agricultural Science*, 6(3), 963-968.
18. Van-Reeuwijk L.P. (1993). Procedures for soil analysis. Wageningen: Technical Paper 9, ISRIC, 16-18.
19. Edori O.S. and Iyama W.A. (2017). Assessment of Physicochemical Parameters of Soils from Selected Abattoirs in Port Harcourt, Rivers State, Nigeria. *Journal of Environmental Analytical Chemistry*, 4(2), 194-201.
20. Ebong G.A., Offiong O.E. and Ekpo B.O. (2014). Seasonal variations in trace metal levels, speciation and physicochemical determinants of metal availability in dumpsite soils within Akwa Ibom State, Nigeria. *Chemistry and Ecology*, 30(5), 403-417.
21. Al-Khashman O.A. and Shawabkeh R.A. (2006). Metals distribution in soils around the cement factory in southern Jordan. *Environmental pollution*, 140(3), 387-394.
22. Fagbote E.O. and Olanipekun E.O. (2010). Speciation of heavy metals in sediment of Agbabu Bitumen deposit area, Nigeria. *Journal of Applied Science and Environmental Management*, 14(4), 47-51.
23. Khan R.S., Kumar N.J. and Patel G.H. (2013). Physicochemical Properties, Heavy Metals Content and Fungal Characterization of an Old Gasoline Contaminated Soil Site in Anand, Gujarat, India. *Environ Exp Biol*, 11, 137-143.
24. Okoye C.O.B. and Agbo K.E. (2011). Dispersion Pattern of Trace Metals in Soils Surrounding Solid Waste Dump in Nsukka. *J. Chem Soc Nigeria*, 36(2), 112-119.
25. Allen S.E., Grimshaw H.M., Parkinson J.A. and Quarmby C. (1974). Chemical analysis of ecological materials. Blackwell Scientific Publications, Oxford, London, UK, 65-70.
26. Rabah A.B., Oyeleke S.B., Manga S.B., Hassan L.G. and Ijah U.J. (2010). Microbiological and Physico-Chemical Assessment of Soil Contaminated with Abattoir Effluents in Sokoto Metropolis, Nigeria. *Science World Journal*, 5(3), 1-4.
27. Cappuccino J.G. and Sherman N. (2002). Techniques for Isolation of Pure Culture. In: *Microbiology: A Laboratory Manual*. 6th ed. Singapore : Pearson Education, Inc., 544.
28. Cheesbrough M. (2006). District Laboratory Practice in Tropical Countries Part 2. UK: Cambridge University Press, 45-70.
29. Brenner D.J., Noel R.K. and Staley J.T. (1982). Bergey's Manual of Systematic Bacteriology. 2nd ed., New York: Springer, 304.
30. Aneja K.R. (2003). Experiments in Microbiology. Plant Pathology and Biotechnology, 4th Ed., New Delhi: New Age International Limited Publishers, 282.
31. Lacatusu R. (2000). Appraising levels of soil contamination and pollution with heavy metals. *European Soil Bureau*, 4, 93-102.
32. Hakanson L. (1980). Ecological Risk Index for Aquatic Pollution Control, A Sedimentological Approach. *Water Research*, 14(8), 975-1001.
33. Rubio B., Nombela M.A. and Vilas F. (2000). Geochemistry of Major and Trace Elements in Sediments of the Ria de Vigo (NW Spain) an Assessment of Metal Pollution. *Marine Pollution Bulletin*, 40(11), 968- 980.
34. Szefer P., Szefer K., Glasby G.P., Pempkowiak J. and Kaliszan R. (1996). Heavy-metal pollution in surficial sediments from the Southern Baltic sea off Poland. *Journal of Environmental Science & Health Part A*, 31(10), 2723-2754.
35. Lu X., Wang L., Lei K., Huang J. and Zhai Y. (2009). Contamination Assessment of Copper, Lead, Zinc, Manganese and Nickel in Street Dust of Baoji, NW China. *Journal of Hazardous Materials*, 161(2-3), 1058-1062.

36. Ho H.H., Swennen R. and Van Damme A. (2010). Distribution and contamination status of heavy metals in estuarine sediments near Cua Ong Harbor, Ha Long Bay, Vietnam. *Geologica belgica*, 13(1-2), 36-47.
37. Tomlinson D.L., Wilson J.G., Harris C.R. and Jeffrey D. W. (1980). Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer meeresuntersuchungen*, 33(1), 566-575.
38. Ediene V.F., Iren O.B. and Idiong M.M. (2016). Effect of Abattoir Effluents on the Physico-chemical Properties of Surrounding Soils in Calabar Metropolis. *International Journal of Advance Research*, 4(8), 37-41.
39. Brady N.C. and Well R.R. (1999). The Nature and Properties of Soil. Pearson Educational Publishers, Delhi, 213.
40. Olayinka O.O., Akande O.O., Bamgbose K. and Adetunji M.T. (2017). Physicochemical characteristics and heavy metal levels in soil samples obtained from selected anthropogenic sites in Abeokuta, Nigeria. *Journal of Applied Sciences and Environmental Management*, 21(5), 883-891.
41. Chibuzor O.J., Nwakonobi T.U. and Itodo I.N. (2017). Influence of Physico-chemical Characteristic of Soils on Heavy Metal Contamination in Makurdi, Benue State. *International Journal of Environmental Science, Toxicology and Food Technology*, 11, 84-92.
42. Franzmeier D.P., Lemme G.D. and Miles R.J. (1995). Organic Carbon in soils of North Central United States. *Soil Science Society America Journal*, 49, 702-708. Rich, K. and Von, I. 2010. Utah State University, USA, Extension and Agriculture Department: Guidelines for Soil Quality, 60.
43. FAO. (1979). Soil Survey investigation for irrigation soils. John Wiley and sons Inc., New York, 23-29.
44. Arias M.E., Gonzalez-Perez J.A. and Ball A.S. (2005). Soil Health: A New Challenge for Microbiologist and Chemists. *Int Microbiol.*, 8, 13-21.
45. Chukwu U.J. and Anuchi S.O. (2016). Impact of Abattoir Wastes on the Physicochemical Properties of Soils within Port Harcourt Metropolis. *The International Journal of Engineering and Science*, 5(6), 17-21.
46. Agbaire P.O. and Emoyan O.O. (2012). Bioaccumulation of heavy metals by earthworm (*Lumbricus terrestris*) and associated soils in domestic dumpsite in Abraka, Delta state, Nigeria. *International journal of plant, animal and environmental sciences*, 2(3), 204-209.
47. Odu C.T.I., Nwoboshi L.C. and Esuruoso O.F. (1985). Environmental Studies (soil and vegetation) of the Nigerian Agip Oil Company Operation Areas. *Proceedings of an international seminar on petroleum industry and the Nigerian environment*, NNPC, Lagos Nigeria, 274-283.
48. Thomson C.J., Marschner H. and Römhild V. (1993). Effect of nitrogen fertilizer form on pH of the bulk soil and rhizosphere, and on the growth, phosphorus, and micronutrient uptake of bean. *Journal of Plant Nutrition*, 16(3), 493-506.
49. Osaki M., Watanabe T. and Tadano T. (1997). Beneficial effect of aluminum on growth of plants adapted to low pH soils. *Soil Science and Plant Nutrition*, 43(3), 551-563.
50. Osakwe S.A. (2016). Contribution of Abattoir Activities in Delta State, Nigeria, to the Soil Properties of their Surrounding Environment. *J Chem Biol Phys Sci.*, 6, 982-991.
51. Akan J.C., Abdulrahman F.I., Sodipo O.A. and Lange A.G. (2010). Physicochemical Parameters in Soil and Vegetable Samples from Gongulon Agricultural Site, Maiduguri, Borno State, Nigeria. *Journal of American Science*, 6(12), 75-87.
52. Onweremadu E.U. (2008). Physico-chemical characterization of a farmland affected by wastewater in relation to heavy metals. *Journal of zhejiang university-science A*, 9(3), 366-372.
53. Chaudhari P.R., Ahire D.V. and Ahire V.D. (2012). Correlation Between Physicochemical Properties and Available Nutrients in Sandy Loam Soils of Haridwar. *Journal of Chemical, Biological and Physical Sciences*, 2 (3), 1493-1500.
54. Nafarnda W.D., Ajayi I.E., Shawulu J.C., Kawe M.S., Omeiza G.K., Sani N.A. and Dantong D.D. (2012). Bacteriological quality of abattoir effluents discharged into water bodies in Abuja, Nigeria. *ISRNVet. Sci.*, 10, 14 -24.
55. WHO/FAO/IAEA. World Health Organization. Switzerland: Geneva. (1996). Trace Elements in Human Nutrition and Health.
56. Adesemoye A.O., Opere B.O. and Makinde S.C.O. (2006). Microbial Content of Abattoir Waste Water and its Contaminated Soil in Lagos, Nigeria. *African Journal of Biotechnology*, 5(20), 1963-1968.
57. Neboh H.A., Iiusanya O.A., Ezekoye C.C. and Orji F.A. (2013). Assessment of Ijebu-Igbo Abattoir Effluent and its Impact on the Ecology of the Receiving Soil and River. *IOSR Journal of Environmental Science and Food Technology*, 7(5), 61-67.
58. Ojo O.F., Adewumi D.G. and Oluwatoyin A.K. (2016). Glutathione-S-transferase production in earthworm (Annelida: Eudrilidae) as a tool for heavy metal pollution assessment in abattoir soil. *Revista de biologia tropical*, 64(2), 779-789.
59. Oyedele D.J., Gasu M.B. and Awotoye O.O. (2008). Changes in Soil Properties and Plant Uptake of Heavy Metals on Selected Municipal Solid Waste Dumpsites in

- Ile-Ife, Nigeria. *African Journal of Environmental Science and Technology*, 2(5), 107-115.
60. Allorge D.O. (1992). Abattoir Design Management and Effluent Disposal in Nigera. University of Ibadan Press. Ibadan, 254.
61. Okoro D., Oviasogie P.O. and Oviasogie F.E. (2011). Soil quality assessment 33 months after crude oil spillage and clean-up. *Chemical Speciation & Bioavailability*, 23(1), 1-6.
62. Iwegbue C.M.A., Egobueze F.E. and Opuene K. (2006). Preliminary assessment of heavy metals levels of soils of an oil field in the Niger Delta, Nigeria. *International Journal of Environmental Science & Technology*, 3(2), 167-172.
63. Hazelton P.A. and Murphy B.W. (2007). Interpreting Soil Test Results: What do all the numbers mean?. *CSIRO Publishing: Melbourne*, 45-67.
64. Rayment G.E. and Higginson F.R. (1992). Cation Exchange Capacity in Australian Laboratory Handbook of Soil and Water Chemical Methods. Inkata Press: Melbourne.
65. Osu C.I. and Okereke V.C. (2015). Heavy Metal Accumulation from Abattoir Wastes on Soils and some Edible Vegetables in Selected Areas in Umuahia metropolis. *International Journal of Current Microbiology and Applied Sciences*, 4(6), 1127-1132.
66. Onianwa P.C. and Essien C.A. (1999). Petroleum hydrocarbon levels in sediments of streams and rivers within Ibadan City, Nigeria. *Bulletin of the Chemical Society of Ethiopia*, 13(1), 83-86.
67. Abubakar G.A. and Tukur A. (2014). Impact of Abattoir Effluent on Soil Chemical Properties in Yola, Adamawa State, Nigeria. *International Journal of Sustainable Agricultural Research*, 1, 100-107.
68. Holland M.D.G., Barton A.D. and Morph S.T. (1989). Land Evaluation for Agricultural Recommendations for Cross River National Park. Oban Division. Prepared by Odwki in Collaboration with INNf.
69. Sumayya B.U., Usman B.U., Aisha U., Shahida A., Mohammad A., Yakubu M.S. and Zainab M. (2013). Determination of Physiochemical Qualities of Abattoir Effluent on Soil and Water in Gandu, Sokoto State. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 4(4), 47-50.
70. Simeon E.O. and Friday K. (2017). Index Models Assessment of Heavy Metal Pollution in Soils within Selected Abattoirs in Port Harcourt, Rivers State, Nigeria. *Singapore Journal of Scientific Research*, 7, 9-15.
71. FEPA (1997). Guidelines and standard for environmental impact assessment in Nigeria. 87-95.
72. Federal Environmental Protection Agency (FEPA) (1999). National Guidelines and Standards for Soil Quality in Nigeria. FEPA, Rivers State Ministry of Environment and Natural Resources, Port Harcourt.
73. Shittu O.S., Ogunwale J.A. and Ayodele O.J. (2010). Total and extractable iron and manganese in soils developed on charnockite in Ekiti State, Nigeria. *International Journal of Water and soil Resources Research*, 1(1-3), 83-93.
74. Hong A.H., Ling L.P. and Selaman O.S. (2014). Environmental Burden of Heavy Metal Contamination Levels in Soil from Sewage Irrigation Area of Geriyo Catchment, Nigeria. *Civil and Environmental Research*, 6(10), 118-124.
75. Atlas R.M. and Bartha R. (2007). Microbial Ecology: Fundamentals and Applications, Benjamin/Cummings Publishing Company Inc, India. 34-39.
76. Alonge D.O. (1991). Textbook of Meat Hygiene in the Tropics. Farm Coe Press, Ibadan, Nigeria, 12-15.