



Physicochemical Properties and Fatty acid composition of Dikanut (*Irvingia Gabonensis*) seed oil

Etong D.I.¹, Mustapha A.O.¹ and Taleat A.A.²

¹Science Laboratory and Technology Department, Federal Polytechnic, P.M.B 420, Offa, Kwara State, NIGERIA

²Science Laboratory And Technology Department, Federal Polytechnic, Ede, Osun State, NIGERIA

Available online at: www.isca.in, www.isca.me

Received 20th November 2014, revised 12nd December 2014, accepted 17th Decmeber 2014

Abstract

Nigeria is one of the countries of the world with a variety of oil seeds, therefore the potential of Nigeria leading Africa, and the world at large, in the area of seeds oil production was examined by evaluating the fat content, physicochemical qualities, and fatty acid content of oil extracted from dikanut (*Irvingia gabonensis*) seed, bought from Owode market in Offa local Government area of Kwara State, Nigerian. The fat content of *Irvingia gabonensis* was 22.50 ± 0.2 %. The chemical qualities iodine, peroxide, saponification, and acid values, free fatty acid, and unsaponifiable matter were analyzed using standard methods, and the values, were found to be $4.50 \pm 0.1 \text{ gI}_2/100\text{g}$, $1.80 \pm 0.1 \text{ meq/kg}$, $187.90 \pm 0.3 \text{ mgKOH/g oil}$, $9.40 \pm 0.2 \text{ mgKOH/g oil}$, $4.70 \pm 0.2\%$ FFA as oleic acid and $1.50 \pm 0.5\%$ respectively. The physical properties were solid at room temperature, grey yellow in colour, melting point ($13 + 1^\circ\text{C}$) and specific gravity $0.88 + 0.2 \text{ kg/dm}^3$. The oil contained six major fatty acid in the following order lauric > myristic > stearic > palmitic > oleic > linolenic acid. The result reveals that the oil extracted can be useful both domestically and industrially.

Keywords: Physicochemical, extraction, *Irvingia gabonensis*, oil, fatty acid.

Introduction

Trees and shrubs with medicinal and nutritional potentials abound in Nigeria. Several of these plants have fruit which have been identified to be nutritionally important¹. In recent times, the desire to conserve resources spent on importation of oil for domestic and industrial use gave renewed impetus in the search for novel sources to complement the traditional ones. Attention has therefore been focused on under-utilized local seeds for possible development and use. There are several of these under-exploited plant seeds in Nigerian².

Dikanut (*Irvingia gabonensis*) belongs to the simarubaceae family and is commonly known as African mango. The fruit is yellow when ripe and the pulp is edible but has a turpentine flavor. The seed ranks as an oil seed and is used as a popular soup thickener in Nigerian³. The Irvingiaceae Kernels are part of the non-timber forest products whose economics and food importance appear undeniable, but despite their importance the Irvingiaceae do not benefit from good valorization and exploitation on a large scale⁴. There are six species of *Irvingia* in the Congo Basin, namely: *Irvingia gabonensis*, *Irvingia grandifolia*, *Irvingia smithii*, *Irvingia wombulu*, *Irvingia excelsa* and *Irvingia robur*⁵, where only *Irvingia gabonensis* and *Irvingia wombulu* species appear to be well known, and are the subject of several transaction and some physiochemical studies^{6,7}.

Many studies undertaken on the nutritional and medicinal value of *Irvingia gabonensis* have been reported⁸⁻¹⁰ etc., work on other

species include *Irvingia smithii*⁴, *Irvingia wombulu*¹¹. Methanolic extract of *Irvingia gabonensis* are used in the treatment of bacterial and fungal infections¹². Seeds oils have extensive demands both for human consumption and for industrial applications¹³ and also have been rated as the second most valuable commodity in the world trade today¹⁴. Numerous researchers^{10,13,15-19}, among others have carried out a lot of analytical works on seeds primarily because of extensive and increasing demands for both human consumption and numerous industrial applications. Consequently, there is an increasing need to perpetually search for oils from non-conventional sources to augment the available ones and also to meet specific applications. Attention has not been focused on under-utilized agricultural foods and industrial products. This study therefore, is designed to search for oil from non-conventional sources because of the increasing needs for oils both for human consumption and

industrial applications. The physicochemical properties and fatty acid content of oil extracted from *Irvingia gabonensis* was studied to a certain its suitability for both domestic and industrial applications.

Material and Methods

Sources and preparation of sample: The sample dikanut (*Irvingia gabonensis*) seed was bought from Owode market in Offa Local government area of Kwara State, Nigeria. The seed was screened to remove the bad ones and stones. They were dried to constant weight in an oven at $60-65^\circ\text{C}$ for 24hours to

reduce moisture content and then grounded using a mechanical grinder, put in an air tight container and stored in a desiccator for further analysis.

Oil extraction and analysis: The seed oil was extracted using the continuous soxhlet extraction technique with petroleum ether (40-60°C) for 5hrs. After removing solvent, using a Rotavapor apparatus, the seed oil obtained was drained under a stream of nitrogen and stored in freezer (-30°C) for subsequent chemical analysis. Standard methods were used in all the analysis saponification value²⁰, unsaponifiable matters, and percentage free fatty acid as oleic acid, and acid valued, peroxide value, iodine value, fat content²¹. Specific gravity was determined using hydrometer²². Refractive index was determined using a refractometer and colour, colorimeter, smoke point, melting and setting points²¹. Fatty acid composition was determined as their methyl esters prepared by boron-trifluoride methanol complex method²¹

Results and Discussion

The studied chemical properties of dikanut (*Irvingia gabonensis*) seed oil are shown in table-1. The fat content was 22.50±0.2%, lower than that reported for castor seeds (57.33±1.63), coconut seeds (42.00±2.21), dikanut (62.80±2.4), groundnut seeds (40.80±0.50), melon seeds, (53.04±1.62), oil beans seeds (52.07±1.45) and palm kernel seeds (54.18±1.88)¹⁰, different varieties of tobacco seeds cultivated in Bangladesh range 42.29±0.60-45.72±1.20¹⁸, seed kernel of *Irvingia smithii* of Congo Basin (54.52±2.61)⁴, *Canarium schwenfurthii* fruits (36.10%), *Balanites aegyptiaca* almonds (48.30%)²³, *Dacryodes edulis* pulp (29-67.50%)²⁴, and *Irvingia gabonensis* seed kernels from two Congo Brazzaville localities, sibiiti (3.69°S; 13.35°E) IGS-(34.55-73.82%), and Ouesso (1.61°N, 16.05°E) IGO-(34.28-62.27%)⁹.

Table-1
Chemical properties of oil extracted from Africa wild mango (*Irvingia gabonensis*) seed

Chemical parameter	Composition
Iodine value (IV) (gI ₂ /100g)	4.50±0.1
Peroxide value (PV) (meq/kg)	1.80±0.1
Saponification value (SV) (mgKOH/g.oil)	187.90±0.3
Acid value (AV) (mgKOH/g oil)	9.40±0.2
% free fatty acid (FFA) as oleic acid	4.70±0.2
Unsaponifiable matter (%)	1.50±0.5
Fat content (%)	22.50±0.2

^a value are means ± standard deviation of triplicate determinations

The saponification value was 187.90±0.30 mgKOH/g oil which was below the range reported for *Irvingia gabonensis* from two localities in Congo Brazzaville (IGO) 189.92±4.6-237.69±72mgKOH/g oil, and (IGS) 196.32±6.2-277.69±2.5 mgKOH g oil⁹. The value was also lower than that reported for *Dacryodes edulis* pulp oil (201)²⁵, those of oil extracted from

conventional oil seeds as soya (189-195), groundnut (187-196) and cotton (189-198)²⁶, it was also higher than that reported for *Chrysophyllum albidum* (126.30±3.50), *Landolphia owariensis* (42.40±3.00) and *Napoleana imperialis* (77.06±4.0) but lower than *Dacryodes edulis* (191.10±3.80) and *Elaeis guineensis* (246.60±4.2)². However, saponification value is used to check adulteration². The relatives high value recorded for this sample is an indication that it have potential for industrial used^{16,25}.

Table-2
Physical properties of oil extracted from African wild Mango (*Irvingia gabonensis*) seed (Means, n = 3)^a

Properties	Value
State at room temperature (27 ± 1°C)	Solid
Colour	Grey yellow
Smoke point (°C)	78±2°C
Melting point (°C)	13 ± 1
Specific gravity Kg/dm ³	0.88 ± 0.2
Setting point (°C)	25.30 ± 1.20

^a values are means ± standard deviation of triplicate determination

Table-3
Fatty acid composition of dikanut (*Irvingia gabonensis*) seed (Mean n=3)^a

Fatty acid	% composition
Oleic	6.90 ± 0.2
Myristic	20.50 ± 0.2
Stearic	11.40 ± 0.1
Linolenic	6.40 ± 0.2
Palmitic	10.30 ± 0.2
Lauric	39.40 ± 0.5

^a value are means ± standard deviation of triplicate determination

Iodine value was 4.50±0.1 gI₂/100g, confirming the values 4.3±0.10-4.80±0.69 for IGO var and 4.10±0.01-4.90±0.05 for IGS var reported for *Irvingia gabonensis* from two localities in Congo Brazzaville⁹. This value was lower compared to that reported for *Cocos nucifera* range (9.93-10.99)^{15,27}, *Colocynthis citrullus* range (110.93-111.46)²⁷. It was higher than 2.65-3.45 reported for *Colocynthis citrullus*, *Cucumeropsis edulis* and *Prunus amygdalus*²⁸, but lower than those reported for most non-conventional plant seeds in Nigeria and Congo Brazzaville^{13,15,29,30}, *Solanum nigrum* L (102.8-103.25), and *Amaranthus hybridus* var. 1 and 2 (109-112.92) from Congo Brazzaville^{29,30} and *Telferia occidentalis* (114.0) and *Citrullus vulgaris* (114.50) from Abeokuta¹⁹. A good drying oil should have iodine value of 180 and above²⁶. Thus, the value obtained for these seed (*Irvingia gabonensis*) classify it as a non-drying oil, as such the seed oil will not be suitable as alky resins for point formulations or used as varnishes, it may however found use in conjunction with amino resins as finishes for certain appliances and in this case, the oil can act as a plasticizers. The

relative low iodine number of the seed oil may be an indicative of the presence of few unsaturated bonds and low susceptibility to oxidative rancidity.

Peroxide value was 1.80 ± 0.1 meq/kg comparable to *C. albidum* (1.8 ± 0.28), *N. Imperialis* (1.55 ± 0.35) but lower than *E. guineensis* (2.12 ± 0.41), *L. owariensis* (2.80 ± 0.50) and *D. edulis* (21.23 ± 1.50)². The value was higher than for *Cocos nucifera* (0.21) but within the range for *Colocynthis citrullus* ($1.53-2.72$)²⁷. The value compares favourably with the range (0.39-7.40) reported for non-conventional seed oils in Nigerian and Congo Brazzaville^{15,19,28,29,30} and the range (0.23-1.80) obtained for the studied conventional oils in the Nigerian markets²⁷. However, it was lower than *B. parii* (97.50), *L. lanceolata* (95.0), *S. setegera* (35.0), *D. Microcarpum* (150.0), *B. Sapida* (135.0) and *S. birrea* (25.0)¹³. Peroxide values are used as indicator for deterioration of oils. Fresh oils have peroxide values less than 10 meq O₂kg⁻¹,²⁶ while values above 20 indicates rancid taste and disagreeable odour. The low peroxide value of this seed oil indicates low level of oxidative rancidity of the oil and also suggests the presence for high level of antioxidant.

Acid value is used as an indicator for edibility of oil and suitability for use in the paint industry. The value for these study was 9.40 ± 0.2 mgKOH/g oil, little above the range 1.20-8.4, mg NaOH g⁻¹ reported for *Telferia occidentalis*, *Citrullus vulgaris*, *Colocynthis citrullus*, (from Akure and Kano), *Pentaclethra macrophylla*, *Treulia africana* and *Cocos nucifera*^{15,19,27,28}, but compared favorably with the range ($3.74 \pm 0.23-25.62 \pm 0.29$) for most conventional oils (Rape seed, soya bean, Wangasa, Kings, Turkey, Filma, Gino and Controller oils) sold in Nigerian markets²⁷. The acid value of the sample studied, fell within the allowable limits for edible oils²⁶, therefore the oil is edible. The free fatty value was $4.70 \pm 0.2\%$ (as oleic acid) was higher than *C. albidum* (1.76 ± 0.10), *D. edulis* (2.78 ± 0.02), *N. impreialis* (2.60 ± 0.07) but lower than *E. guineensis* (7.06 ± 0.01) and *L. owariensis* (7.70 ± 0.11)², also higher than that reported for three varieties of tobacco seed oil ($1.51-2.30$)¹⁸. The oil has low acid value and a corresponding low level of free fatty acid, which suggests low level of hydrolytic and lipolytic activities in the oil, thus the seed oil studied could be a good source of raw materials for industries.

The unsaponifiable matter was $1.50 \pm 0.50\%$, which was comparable to that reported for three tobacco varieties 1.39-1.45%¹⁸, but lower than that obtained for *Psophocarpus tetragonolobus* (1.63%) and orchid fruits *Myristica* (1.79%) oil seeds³¹. Unsaponifiable matter includes hydrocarbons, higher alcohols and sterols (e.g. Cholesterol, phytosterol)¹⁸. Most oils and fats of normal purity contain less than 2% of unsaponifiable matter.

Table II showed the physical properties of oil extracted from dikanut (*Irvingia gabonensis*) seeds. The oil was solid at room temperature ($27 \pm 1^\circ\text{C}$) and grey yellow in colour. The specific

gravity was 0.88 ± 0.2 , comparable to the range 0.88-0.93 for *Cocos nucifera* seeds but little below the range for *Cocoynthia citrullus* (0.90-0.92)²⁷. The specific gravity value 0.88 ± 0.2 , ranked favorably with the range of specific gravities 0.81-0.928 reported for other oils, fats and waxes from common plant seeds in Nigerian^{2, 15, 19}. However, the value was low compared to those reported for seeds from Akure, *Colocynthis citrullus* (1.51), *Cucumeropsis, edulis* (1.67) and *Prunus amygdalus* (1.71)²⁸. The smoke point was ($78.0 \pm 0.2^\circ\text{C}$), lower than that reported for three varieties of tobacco seeds oil ($214 \pm 1.41-230 \pm 0.81$)¹⁸, soyabean oil ($250.00 \pm 5.77^\circ\text{C}$), *Blighia sapida* ($220 \pm 2.88^\circ\text{C}$)³². Smoke point of a fatty material is measure of its thermal stability when heated in contact with the air. Fatty acids are much less stable than glycerides, hence the smoke points of ordinary oils depend principally upon their content of free fatty acids. The melting point was $13 \pm 1.0^\circ\text{C}$, significantly lower than range reported for selected Nigerian oil seed ($34.0 \pm 3.5-72.0 \pm 2.6^\circ\text{C}$)¹⁰. Since melting point decrease with increase in the degree of unsaturation¹⁰, the lower melting point oils may be useful in the manufacture of soft and easy-to- digest margarine and have be shown to be valuable in the manufacture of oil creams. The setting point was $25.30 \pm 1.20^\circ\text{C}$ which was within the range reported for some selected Nigerian oil seeds ($4.67 \pm 0.30-26.30 \pm 1.50^\circ\text{C}$)¹⁰.

Table-3, presents the fatty acid composition of dikanut (*Irvingia gabonensis*). The following fatty acids were found in the following percentage lauric acid (39.40 ± 0.5), myristic acid (20.50 ± 0.3), stearic acid (11.40 ± 0.1), palmitic acid (10.30 ± 0.2), oleic acid (6.90 ± 0.2) and linolenic acid (6.40 ± 0.2). The sample was richer than *Citrullus lanatus* seed-linolenic acid (5.29), stearic acid (8.33), but comparable in palmitic acid (10.57), and lower in terms oleic acid (13.65)¹⁷. Several authors reported that *Irvingia gabonensis* seed kernels oil may be regarded as myristic-lauric oil because myristic acid was most abundant, followed by lauric acid⁷, but in this study the reverse was the case lauric was the most abundant followed myristic, thus, it may be regarded as lauric- myristic oil.

Conclusion

Taken as a whole, in this study, the physicochemical characteristics can be helpful to identify the quality of oil and oil products for possible industrial or commercial uses. The oil extracted, is saturated, a source of lauric and myristic acid, which is the characteristics of vegetable oils. From the quality point of view, *Irvingia gabonensis* seed oil classified as lauric-myristic oils, is comparable to other oils and can be utilized for both industrial and domestic purpose. The oil has a considerable high quality and good shelf life.

Reference

1. Ihekoronye A.I. and Ngoddy P.O., Integrated Food Science and Technology for the tropics, Macmillan Publishers, 82-93 (1985)

2. Akubugwo I.E. and Ugbugu A.E., Physicochemical studies on oils from five selected Nigerian plant seeds. *Pak. Journal of Nutrition*. **6 (10)**: 75-78 (2007)
3. Onyeike E.N., Olungwe T. and Uwakwe A.A., Effect of heat treatment and defatting on proximate composition of some Nigerian local soup thickeners, **53**, 173- 175 (1995)
4. Loumouamou B.M., Gomoufatan J.P.M., Silou T., Nzikou J.M., Mbaya Gindo G.V., Figueredo G. and Chalard J.P., Extraction and Chemical composition of seed kernel oil from *Irvingia smithii* of Congo Basin, *Advance Journal of Food Sci. and Tech.*, **5(5)**, 506-513 (2013)
5. Makita-madzou J.P., Report of Congo IPGRI. Program of forest genetic resource in Africa South of the Sahara, Eyog Matig, O., O. Gande Gaoue and B. Dossou (Eds), Food Tree Species Network (French), Minutes of the first meeting of the network, CNSF Ouagadougou, Burkina Faso, 11-13, (2000)
6. Silou T.H., Biyoko S., Heron S., Tchaplala A. and Maloumbi M.G., Physicochemical characteristics and technological potentialities of *Irvingia gabonensis* seeds, *Riv. Ital Sostanze, Gr.*, **81**, 49-57 (2004)
7. Matos L., Nzikou J.M., Kimbonguila A., Ndangui C.B., Pambou-Tobi N.P.G., Abena A.A., Silou T.H., Scher J. and Desobry S., Composition and nutritional properties of seeds and oils from *Terminalia catappa* L. *Advance Journal of Fod Sci. and Tech.*, **1(1)**, 71-76 (2009a)
8. Ngondi J., Oben J. and Minka S., The effect of *Irvingia gabonensis* seeds on body weight and blood lipids of obese subject in Cameroon, *Lipids Health Diseases*, **4**, 12-17 (2005)
9. Matos, L., Nzikou, J.M., Matoubua E., Pandzou-Yembe, V.N., Guembot Mapepoulou, T., Linder, M. and Desobry, S., Studies of *Irvingia gabonensis* seed kernels: Oil technological applications, *Pak. Jour. of Nutri.*, **8(2)**, 151-157 (2009b)
10. Onyeike, E.N. and Acheru, G.N., Chemical composition of selected Nigerian oil seeds and physicochemical properties of the oil extracted, *Food Chem.* **77**, 431-437 (2002)
11. Silou, T., Loumouamou, B.W., Nsikabak, S., Kinkela, T., Nzikou, J.M., Chalard, P. and Figueredo, G., Contribution to the varietal delimitation of *Irvingia gabonensis* and *Irvingia woubulu* chemical composition variability of fats extracted from kernels, *Journ. Food. Tech.*, **9(1)**, 36-42 (2011)
12. Kuete, V., Wabo, G.F., Ngameni, B., Tsafack Mbaveng, A., Metuno, R., Penlap Beng, V., Marion Meyer, J.J. and Lall, N., Antimicrobial activities of the methanolic extract, fractions and compounds from the stem bark of *Irvingia gabonensis* (Ixonanthaceae), *J. Ethnopharmacology*, **114**, 54-60 (2007)
13. Kyari, M.Z., Extraction and characterization of seed oils, *Int. Agrophysics*, **22**, 139-142 (2008)
14. Ige, M.M., Ogunsua, A.O. and Oke, L.O., Functional properties of proteins of some Nigerian oil seeds : Conophor seeds and three varieties of melon seeds, *Journal Agric. Food Chem.*, **32**, 822-825 (1984)
15. Akubugwo, I.E., Chinyere , G.C, and Agbogou, A. E., Comparative studies on oils from some common plants seeds in Nigerian. *Pak. Journal of Nutrition.*, **7**, 570-573 (2008)
16. Olagunju E.O., Extraction and characteristic of vegetable oil from Mango seed, *Mangifera indica*. *IOSIR Journal of Applied chemistry*, **5(3)**, 6-8 (2013)
17. Edidiong A.E. and Ubong M.E., Analysis of *Citrullus lanatus* seed oil obtained from Southern Nigerian., *Elixir Org. Chem.*, **54**, 12700-12703 (2013)
18. Abbas M.A., Abu M.S., Rajesh K.R., Sarmina Y. and Astaq, M.K., Comparative study on characteristics of seed oils and nutritional composition of seeds from different varieties of tobacco (*Nicotiana tabacum* L.) cultivated in Bangladesh, *Asian Journal of Biochemistry*, **3(4)**, 203-212 (2008)
19. Yusuf A.A., Adewuyi S. and Lasisi A.A., Physicochemical composition of leaves, meals and oils of Fluted pumpkin (*Telfairia Occidentalis*) and melon (*Citrullus vulgaris*), *Agric. Journal*, **1**, 32-35 (2006)
20. Pearson D., The chemical analysis of food, church hill London., 580-584 (1981)
21. AOAC, Official methods of analysis, 17th eds. Washington D.C: Association of official analytical chemists, (1997)
22. Gerpen V.J., Shanks B., Pruszko R., Clement D. and Knothe G., Biodiesel analytical methods. National Renewable Energy Laboratory, Midwest Research Institute, Battelle (2004)
23. Nzikou J.M., Mvoula Tsier M., Matouba E., Ouamba J.M., Kapseu C., Parmentier M, and Desobry S., A study on gumbo seed grown in Congo Brazzaville for its food and Industrial applications, *Afri. Journ. Biotechn.*, **5**, 2469-2475 (2006)
24. Silou T., Massamba D., Goma Maniongui J., Maloumbi G. and Biyoko S., Post-harvest losses by natural softening of safou pulp (*Dacryodes edulis*) in Congo-Brazzaville, *Journ. Food Engineering*, (Article in Press) (2006)
25. Nzikou J.M., Mvoula-Tsier M., Matos L., Matouba E., Ngakegni-Limbili A.C., Linder M. and Desobry S., *Solanum nigrum* L. Seeds as an alternative sources of edible lipids and nutrients in Congo Brazzaville, *Journ. Appl. Sci.*, **7**, 1107-1115 (2007)

26. Alimentarius Codex Commission Recommended International Standard for edible fats and oils, Edn 1 vol 11, (1) FAO/WHO, Rome, 1-179 (1982)
27. Obasi N.A., Ukadilonu J., Eze E., Akubugwo E.I. and Okorie U.C., Proximate composition, Extraction, Characterization and comparative Assessment of coconut (*Cocos nucifera*) and melon (*Colocynthis citrullus*) seeds and seed oils, *Pak. Journal of Biological Sci.*, **15(1)**, 1-9 (2012)
28. Akpambang V.O.E.; Amoo I.A. and Izuagie O., Comparative compositional analysis on two varieties of melon (*Colocynthis citrullus* and *Cucumeropsis*) and a variety of almond (*Prunus amygdalus*), *Res. Journ. Agric. Bio. Sci.*, **2(3)**, 20-25 (2008)
29. Dhellot J.R., Matouba E., Maloumbi M.G, Nzikou J.M. and Ngoma D.G.S., Extraction, Chemical composition and nutritional characterization of vegetable oils : Case of *Amaranthus hybridus* (Var 1 and 2) of Congo. *African J. Biotech.*, **5**, 1095-1101 (2006a)
30. Dhellot J.R., Matouba E., Maloumbi M.G., Nzikou J.M. and Dzondo M.G., Extraction and nutritional properties of *Solanum nigrum* L. seed oil., *Afr. J. Biotechnology*, **5**, 987-991 (2006b)
31. Amoo I.A., Adebayo O.T. and Oyeleye A.O., Chemical Evaluation of Winged beans (*Psophocarpus tetragonolobus*), pitanga cherrites (*Eugenia uniflora*) and orchid fruit (*Myristica*), *Afri. J. Food Agric. Nutri. Develop.*, **6(2)**, 1-12 (2006)
32. Oladiji A.T., Shoremekun K.L. and Yakubu M.T.Y., Physicochemical Properties of the oil from the fruit of *Blighia sapida* and toxicological evaluation of the oil-based diet in Wistar rats, *Journal of medicinal food*, **12(5)**, 1127-1135 (2009)